

Comprehensive Exam – Linear Algebra
June 2008

1. Determine if each of the following statements is TRUE or FALSE by giving a short proof or a counterexample. Throughout, $A \in \mathcal{M}_n(\mathbb{C})$, unless otherwise indicated.

- (a) If all the eigenvalues of A are zero, then $A = 0$.
- (b) If λ is an eigenvalue of A and $\mu \in \mathbb{C}$, then $\lambda - \mu$ is an eigenvalue of $A - \mu I$.
- (c) If A is diagonalizable and its eigenvalues are equal, then A is diagonal.
- (d) If λ is an eigenvalue of A and A is nonsingular, then λ^{-1} is an eigenvalue of A^{-1} .

2. Let V be a vector space over the field \mathbb{F} , and let T be a linear map from V to \mathbb{F} . Prove that if $u \in V$ is not in $N(T)$, the nullspace of T , then

$$V = \{au : a \in \mathbb{F}\} \oplus N(T).$$

3. If V is a vector space and U, T two linear transformations on V which commute, $UT = TU$, prove that any eigenspace of T is U -invariant.

4. Given an orthogonal operator U on a real inner product space V , prove that

- (a) Any eigenvalues of U have absolute value 1.
- (b) If the $\dim V$ is odd, then U has at least one real eigenvalue.
- (c) Give an example ($\dim V$ even) when an orthogonal operator does not have any eigenvalues.

5. Prove that if there exist a linear map on V whose null space $N(T)$ and range $R(T)$ are both finite dimensional, then V is finite dimensional.

6. Let $A = \begin{pmatrix} 1 & 1 & 1 & 0 & 0 \\ 0 & 2 & 1 & 3 & 0 \\ 0 & 0 & 2 & 0 & 0 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix} \in \mathcal{M}_5(\mathbb{R})$.

- (a) Compute the eigenvalues of A and both the characteristic and minimal polynomial for A .
- (b) Find the Jordan form for A .

[Recall that the largest integer d such that $(t - \lambda)^d$ divides the minimal polynomial of a matrix equals the largest size of a Jordan block corresponding to the eigenvalue λ appearing in the Jordan canonical form of that matrix.]