

# SOLUTIONS

## EXAM I – Linear Algebra

• **Problem 1**

Let  $V$  be the subspace of  $\mathbb{R}^4$  generated by the vectors  $v_1 = (1, 1, 0, 0)$ ,  $v_2 = (0, 1, 1, 0)$ ,  $v_3 = (0, 0, 1, 1)$  and  $W$  generated by the vectors  $w_1 = (1, 0, 1, 0)$ ,  $w_2 = (0, 2, 1, 1)$ ,  $w_3 = (1, 2, 1, 2)$  in  $\mathbb{R}^4$ .

(a) Determine the dimensions of  $V$  and  $W$ .

**Solution.**  $\dim(V) = 3$  since  $V = \text{span}\{v_1, v_2, v_3\}$  and the vectors  $v_1, v_2, v_3$  are lin. independent. Indeed,  $a_1v_1 + a_2v_2 + a_3v_3 = 0 \Rightarrow a_1(1, 1, 0, 0) + a_2(0, 1, 1, 0) + a_3(0, 0, 1, 1) = (0, 0, 0, 0) \Rightarrow (a_1, a_1 + a_2, a_2 + a_3, a_3) = (0, 0, 0, 0) \Rightarrow a_1 = 0, a_2 = 0, a_3 = 0$ .

Same argument for  $W$ :  $\dim(W) = 3$  since  $w_1, w_2, w_3$  are lin. ind. (a similar argument is needed!)

(b) Find a basis for the sum  $V + W$ .

**Solution.** Since  $\{v_1, v_2, v_3\}$  is lin. ind. and  $w_1 \notin \text{span}\{v_1, v_2, v_3\}$  (This needs proof!) then, by Theorem 1.7,  $\{v_1, v_2, v_3, w_1\}$  is lin. ind. in  $\mathbb{R}^4$ . Hence  $\text{span}\{v_1, v_2, v_3, w_1\} = \mathbb{R}^4$ . Now  $\{v_1, v_2, v_3, w_1\} \subset V + W$  means that  $V + W = \mathbb{R}^4$ . We conclude that  $\{v_1, v_2, v_3, w_1\}$  is a basis for  $V + W$ .

(c) Find a basis for the intersection  $V \cap W$ . Verify that  $\dim(V + W) + \dim(V \cap W) = \dim(V) + \dim(W)$ .

**Solution.** For a vector  $v$  to belong to  $V \cap W$ , it is necessary and sufficient to have  $v$  as a linear combination of both bases  $\{v_1, v_2, v_3\}$  (for  $V$ ) and  $\{w_1, w_2, w_3\}$  for  $W$ :

$$v = a_1v_1 + a_2v_2 + a_3v_3 = b_1w_1 + b_2w_2 + b_3w_3, \quad \text{for } a_i, b_i \in \mathbb{R}, i = 1, 2, 3.$$

Hence  $a$ 's and  $b$ 's are solutions to the underdetermined linear system

$$a_1 \begin{pmatrix} 1 \\ 1 \\ 0 \\ 0 \end{pmatrix} + a_2 \begin{pmatrix} 0 \\ 1 \\ 1 \\ 0 \end{pmatrix} + a_3 \begin{pmatrix} 0 \\ 0 \\ 1 \\ 1 \end{pmatrix} = b_1 \begin{pmatrix} 1 \\ 0 \\ 1 \\ 0 \end{pmatrix} + b_2 \begin{pmatrix} 0 \\ 2 \\ 1 \\ 1 \end{pmatrix} + b_3 \begin{pmatrix} 1 \\ 2 \\ 1 \\ 2 \end{pmatrix}$$

Performing reduced row echelon form on the matrix  $\begin{pmatrix} 1 & 0 & 0 & 1 & 0 & 1 \\ 1 & 1 & 0 & 0 & 2 & 2 \\ 0 & 1 & 1 & 1 & 1 & 1 \\ 0 & 0 & 1 & 0 & 1 & 2 \end{pmatrix}$  yields:  $\begin{pmatrix} 1 & 0 & 0 & 0 & 1 & 2 \\ 0 & 1 & 0 & 0 & 1 & 0 \\ 0 & 0 & 1 & 0 & 1 & 2 \\ 0 & 0 & 0 & 1 & -1 & -1 \end{pmatrix}$

hence the system above is equivalent to

$$a_1 \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix} + a_2 \begin{pmatrix} 0 \\ 1 \\ 0 \\ 0 \end{pmatrix} + a_3 \begin{pmatrix} 0 \\ 0 \\ 1 \\ 0 \end{pmatrix} = b_1 \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix} + b_2 \begin{pmatrix} 1 \\ 1 \\ 1 \\ -1 \end{pmatrix} + b_3 \begin{pmatrix} 2 \\ 0 \\ 2 \\ -1 \end{pmatrix},$$

or  $a_1 = b_2 + 2b_3, a_2 = b_2, a_3 = b_2 + 2b_3, 0 = b_1 - b_2 - b_3$ . As a consequence, we can choose arbitrarily  $b_2 = s$  and  $b_3 = t$ , then  $a_1 = s + 2t, a_2 = s, a_3 = s + 2t, b_1 = s + t$ .

Then every  $v \in V \cap W$  can be expressed as  $v = (s + 2t)v_1 + sv_2 + (s + 2t)v_3 = (s + t)w_1 + sw_2 + tw_3$ . or  $v = s(v_1 + v_2 + v_3) + t(2v_1 + 2v_3) = s(w_1 + w_2) + t(w_1 + w_3)$ . It means that a basis for  $V \cap W$  consists of the two vectors  $v_1 + v_2 + v_3 = w_1 + w_2 = (1, 2, 2, 1)$  and  $2v_1 + 2v_3 = w_1 + w_3 = (2, 2, 2, 2)$ .

One verifies that  $\dim(V + W) + \dim(V \cap W) = 4 + 2 = 6 = 3 + 3 = \dim(V) + \dim(W)$ . Q.E.D.

• **Problem 2**

Given a vector space  $V$ , show that when two finite dimensional subspaces  $W_1$  and  $W_2$  satisfy

$$\dim(W_1 + W_2) = \dim(W_1 \cap W_2) + 1$$

then either  $W_1 \subset W_2$  or  $W_2 \subset W_1$  and  $|\dim(W_1) - \dim(W_2)| = 1$ .

**Solution.** Denote  $n = \dim(W_1 + W_2)$ . We distinguish two cases:

Case 1. If  $W_1 \subseteq W_2$  then obviously  $W_1 \cap W_2 = W_1$  and  $W_1 + W_2 = W_2$ ,  $\dim(W_1) = n - 1$  and  $\dim(W_2) = n$ .

Case 2. If  $W_1 \not\subseteq W_2$ , then there exists  $w \in W_1 \setminus W_2$ . Let  $\{v_1, v_2, \dots, v_{n-1}\}$  be a basis for  $W_1 \cap W_2$ . We claim that  $\{v_1, v_2, \dots, v_{n-1}, w\}$  is then a basis in  $W_1 + W_2$ . Indeed, by Theorem 1.7,  $\{v_1, v_2, \dots, v_{n-1}, w\}$  is lin. independent and it contains exactly  $n$  elements, therefore it generates (is a basis for)  $W_1 + W_2$ . This means  $W_1 = W_1 + W_2$ ,  $\dim(W_1) = n$  and  $\dim(W_2) = n - 1$ . Q.E.D.

• **Problem 3**

Consider a linear transformation  $T : V \rightarrow V$  over a finite dimensional vector space  $V$  with  $\dim(V) = n$ . Let  $v \in V$  be a given vector such that  $T^n(v) = 0$ , but  $T^{n-1}(v) \neq 0$ .

(a) Prove that the set  $\beta = \{v, T(v), T^2(v), \dots, T^{n-1}(v)\}$  forms a basis for  $V$ .

**Solution.** Since the set contains exactly  $n = \dim(V)$  elements, to show it is a basis for  $V$  it is sufficient to show that it is a linear independent set in  $V$ . Assume

$$a_0 v + a_1 T(v) + a_2 T^2(v) + \dots + a_{n-1} T^{n-1}(v) = 0.$$

Applying  $T^{n-1}$  to both sides and using linearity and the fact that  $T^n(v) = 0$ , we obtain  $a_0 T^{n-1}(v) = 0$ , and since  $T^{n-1}(v) \neq 0$ , we conclude  $a_0 = 0$ . Hence

$$a_1 T(v) + a_2 T^2(v) + \dots + a_{n-1} T^{n-1}(v) = 0.$$

We continue by applying  $T^{n-2}$  to both sides, to obtain  $a_1 T^{n-1}(v) = 0$ , hence  $a_1 = 0$ , etc. After  $n$  iterations of this procedure, we obtain  $a_0 = a_1 = a_2 = \dots = a_{n-1} = 0$ . Q.E.D.

(b) Find a basis for the null-space  $N(T)$  and the range  $R(T)$  of  $T$  and verify the dimension theorem.

**Solution.** Since  $T^n(v) = 0$  it means  $T^{n-1}(v) \in N(T)$ . Also, it is clear that  $\{T(v), T^2(v), \dots, T^{n-1}(v)\} \subset R(T)$ . Based on these observations, we claim that

$$N(T) = \text{span}\{T^{n-1}(v)\} \text{ and } R(T) = \text{span}\{T(v), T^2(v), \dots, T^{n-1}(v)\}$$

Indeed, by the inclusions above it means that  $\dim(N(T)) \geq 1$  and  $\dim(R(T)) \geq n - 1$ . But the dimension theorem guarantees that  $\dim(N(T)) + \dim(R(T)) = n$ , hence  $\dim(N(T)) = 1$  and  $\dim(R(T)) = n - 1$ . Thus the claim is proved.

(c) Calculate  $[T]_\beta$ .

**Solution.** It is easy to construct the matrix representation of  $T$  in the basis  $\beta$ , since  $T(v) = 0 \cdot v + 1 \cdot T(v) + \dots + 0 \cdot T^{n-1}(v)$  etc. We obtain:

$$[T]_\beta = \begin{pmatrix} 0 & 0 & 0 & \dots & 0 & 0 \\ 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 1 & 0 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & 0 & 0 \\ 0 & 0 & 0 & \dots & 1 & 0 \end{pmatrix}$$

• **Problem 4**

Let  $V$  be a finite dimensional vector space.

(a) Show that for any given subspace  $W_1$  of  $V$ , there exists a subspace  $W_2$  of  $V$  such that  $W_1 \oplus W_2 = V$

**Solution.** Since  $V$  is finite dimensional, say  $\dim(V) = n$ , so is  $W_1$ . Denote  $\dim(W_1) = k$ . Let  $\beta_1 = \{v_1, v_2, \dots, v_k\}$  be a basis for  $W_1$ . Then we can complete  $\beta_1$  to a basis  $\beta = \{v_1, v_2, \dots, v_k, v_{k+1}, \dots, v_n\}$  for  $V$ . We claim that  $W_2 = \text{span}\{v_{k+1}, \dots, v_n\}$  is a subspace in  $V$  satisfying  $W_1 \oplus W_2 = V$ . Indeed, one easily verifies that  $W_1 + W_2 = V$  and  $W_1 \cap W_2 = \{0\}$ , using the properties of the basis  $\beta$ .

(b) Let  $T : V \rightarrow V$  be the projection on  $W_1$  along  $W_2$ , where  $W_1$  and  $W_2$  are as in part (a). Show that  $T^2 = T$ .

**Solution.** Let  $v \in V = W_1 \oplus W_2$ , with the unique representation  $v = v_1 + v_2$ , with  $v_1 \in W_1, v_2 \in W_2$ . Then  $T^2(v) = T(T(v)) = T(v_1) = v_1$ , the last equality since  $v_1 = v_1 + 0$  is the unique representation of  $v_1 \in W_1$  as a sum of elements in  $W_1$  and  $W_2$ . Hence  $T^2(v) = T(v)$ . Since  $v$  was arbitrary in  $V$ , it follows that  $T^2 = T$ .

(c) Conversely, show that any linear transformation  $T : V \rightarrow V$  satisfying  $T^2 = T$  is a projection on  $R(T)$  along  $N(T)$ .

**Solution.**

Consider  $T$  a linear transformation satisfying  $T^2 = T$ . Denote  $W_1 = R(T)$  and  $W_2 = N(T)$ . We claim that  $N(T) \oplus R(T) = V$  and  $T$  is the projection onto  $W_1$  along  $W_2$ .

Indeed, for  $v \in V$ , write  $v = T(v) + v - T(v)$ . We have  $T(v) \in R(T)$  and  $v - T(v) \in N(T)$ , since  $T(v - T(v)) = T(v) - T^2(v) = T(v) - T(v) = 0$ . Hence  $V = W_1 + W_2$ .

To show  $W_1 \cap W_2 = \{0\}$ , let  $v \in W_1 \cap W_2$ . It implies that  $v = T(w)$  for some  $w \in V$  and  $T(v) = 0$ . Then  $0 = T(v) = T^2(w) = T(w) = v$ , hence  $v = 0$ .

Using the unique representation of any  $v$  in  $W_1 \oplus W_2$  as  $v = T(v) + (v - T(v))$ , it is clear that  $T(v) = T(v)$  implies that  $T$  acts on the direct sum as the projection on  $W_1$  along  $W_2$ .

• **Problem 5**

Given a  $2 \times 2$  matrix  $P = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ , define the map  $T : \mathcal{M}_{2 \times 2}(\mathbb{R}) \rightarrow \mathcal{M}_{2 \times 2}(\mathbb{R})$  given by left multiplication by  $P$ :

$$T(A) = PA, \quad \text{for } A \in \mathcal{M}_{2 \times 2}(\mathbb{R}).$$

(a) Show that  $T$  is linear.

**Solution.** We easily verify  $T(A_1 + A_2) = P(A_1 + A_2) = PA_1 + PA_2 = T(A_1) + T(A_2)$  and  $T(cA) = PcA = cPA = cT(A)$  for  $c \in \mathbb{R}$ , by using the properties of matrix addition and multiplication.

(b) Determine the matrix representation of  $T$  in the standard ordered basis for  $\mathcal{M}_{2 \times 2}(\mathbb{R})$

$$\left\{ \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}, \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} \right\}.$$

**Solution.** We compute  $\begin{pmatrix} a & b \\ c & d \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} = \begin{pmatrix} a & 0 \\ c & 0 \end{pmatrix}$  etc and therefore obtain

$$[T] = \begin{pmatrix} a & 0 & b & 0 \\ 0 & a & 0 & b \\ c & 0 & d & 0 \\ 0 & c & 0 & d \end{pmatrix}.$$

• **Problem 6**

Let  $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$  be defined as the reflection about the plane  $x + y + 2z = 0$ .

(a) Explain in a few words why  $T$  is a linear transformation.

**Solution.** Addition of vectors obeys the parallelogram rule, hence whether it is performed first between two vectors  $v, w$  and then reflected about the plane, or performed directly on the mirror of  $v$  and  $w$  yields the

same result. Same with scalar multiplication.

(b) Find an expression for  $T(a, b, c)$ , for any  $(a, b, c) \in \mathbb{R}^3$ .

[Hint: Find a convenient basis  $\beta'$  in which  $[T]_{\beta'}$  is easy to be computed, then use the change of bases formula to compute  $[T]_{\beta}$ , where  $\beta$  is the standard ordered basis in  $\mathbb{R}^3$ .]

**Solution.** We pick a convenient basis (with 3 vectors) in  $\mathbb{R}^3$  so that two vectors belong to the plane  $x + y + 2z = 0$ , say  $v_1 = (2, 0, -1)$  and  $v_2 = (0, 2, -1)$  and a third vector is perpendicular to the plane, say  $v_3 = (1, 1, 2)$ . Then  $T(v_1) = v_1, T(v_2) = v_2$ , and  $T(v_3) = -v_3$ . Therefore, in the ordered basis  $\beta' = \{v_1, v_2, v_3\}$ ,  $T$  is represented by the matrix:

$$[T]_{\beta'} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$$

Denoting my  $\beta$  the standard ordered basis in  $\mathbb{R}^3$ , we know the change of bases matrix from  $\beta$  to  $\beta'$  is  $Q = \begin{pmatrix} 2 & 0 & 1 \\ 0 & 2 & 1 \\ -1 & -1 & 2 \end{pmatrix}$ . We compute  $Q^{-1} = \frac{1}{12} \begin{pmatrix} 5 & -1 & -2 \\ -1 & 5 & -2 \\ 2 & 2 & 4 \end{pmatrix}$  use the relation  $[T]_{\beta'} = Q^{-1}[T]_{\beta}Q$  to obtain

$$[T]_{\beta} = Q[T]_{\beta'}Q^{-1} = \frac{1}{3} \begin{pmatrix} 2 & -1 & -2 \\ -1 & 2 & -2 \\ -2 & -2 & -1 \end{pmatrix}.$$

We conclude that  $T(a, b, c) = \frac{1}{3}(2a - b - 2c, -a + 2b - 2c, -2a - 2b - c)$  is the expression for this transformation (in the standard coordinate system).

• **Problem 7\***

(a) Given a vector space  $V$ , show that the union of an increasing sequence of subspaces of  $V$  is a subspace of  $V$ .

**Solution.** Let  $W_1 \subseteq W_2 \subseteq W_3 \subseteq \dots \subseteq W_k \subseteq W_{k+1} \subseteq \dots$  be an increasing sequence of subspaces of  $V$ . Denote  $W = \bigcup_{k \geq 1} W_k$ . We show  $W$  is a subspace by showing that it is closed under addition and scalar multiplication. Indeed, let  $c$  be a scalar,  $u, v \in W$ , then there exists  $k \geq 1$  such that  $u, v \in W_k$ . But since  $W_k$  is subspace,  $u + cv \in W_k$  as well, hence  $u + cv \in W$ .

(b) Let  $V$  be the vector space of infinite sequences of real numbers that converge to 0 and  $T$  be the left shift operator on  $V$  (see exercise 21, page 76). Show that  $W$ , the subset of sequences that have only finitely many nonzero terms, satisfies

$$W = \bigcup_{k \geq 1} N(T^k)$$

and deduce that  $W$  is a subspace of  $V$ , which is also invariant under  $T$ . Determine a basis for  $W$ .

**Solution.** By the definition of  $T$ ,  $N(T)$  = set of all sequences that have all terms zero (except possibly the first term):

$$N(T) = \{\mathbf{a} = (a_1, 0, 0, \dots) \mid a_1 \in \mathbb{R}\}.$$

Similarly,

$$N(T^2) = \{\mathbf{a} = (a_1, a_2, 0, \dots) \mid a_1, a_2 \in \mathbb{R}\}.$$

and more generally

$$N(T^k) = \{\mathbf{a} = (a_j) \mid a_j \in \mathbb{R}, a_j = 0 \text{ whenever } j > k\}.$$

Clearly  $N(T) \subset N(T^2) \subset \dots$  (This is true in fact for ANY linear transformation  $T : V \rightarrow V$ ), and, moreover  $W$ , being the set of all sequences that have only finitely many nonzero terms, equals their union. By part (a),  $W$  is a subspace of  $V$ .

$W$  is invariant under  $T$  since the action of  $T$  does not increase the number of nonzero terms in a sequence (on the contrary, it may decrease the number of nonzero terms.) More precisely,

$$T(N(T^k)) = N(T^{k-1}), \text{ for all } k, \Rightarrow T(W) = T\left(\bigcup_{k \geq 1} N(T^k)\right) = \bigcup_{k \geq 1} N(T^{k-1}) = W.$$

A basis for  $W$  consists of the infinite set  $\{e_1, e_2, \dots\}$ , where  $e_k$  is the sequence that has zero everywhere except the  $k^{\text{th}}$  term, which is equal to 1: e.g.  $e_1 = (1, 0, 0, \dots), e_2 = (0, 1, 0, \dots)$ . (Show this is basis for  $W$ !)