

Isolation and Characterization of Viruses Related to the SARS Coronavirus from Animals in Southern China

Y. Guan,^{1*} B. J. Zheng,^{1*} Y. Q. He,² X. L. Liu,² Z. X. Zhuang,² C. L. Cheung,¹ S. W. Luo,¹ P. H. Li,¹ L. J. Zhang,¹ Y. J. Guan,¹ K. M. Butt,¹ K. L. Wong,¹ K. W. Chan,³ W. Lim,⁴ K. F. Shortridge,¹ K. Y. Yuen,¹ J. S. M. Peiris,¹ L. L. M. Poon¹

A novel coronavirus (SCoV) is the etiological agent of severe acute respiratory syndrome (SARS). SCoV-like viruses were isolated from Himalayan palm civets found in a live-animal market in Guangdong, China. Evidence of virus infection was also detected in other animals (including a raccoon dog, *Nyctereutes procyonoides*) and in humans working at the same market. All the animal isolates retain a 29-nucleotide sequence that is not found in most human isolates. The detection of SCoV-like viruses in small, live wild mammals in a retail market indicates a route of interspecies transmission, although the natural reservoir is not known.

Severe acute respiratory syndrome (SARS) recently emerged as a human disease associated with pneumonia (1). This disease was first recognized in Guangdong Province, China, in November 2002. Subsequent to its introduction to Hong Kong in mid-February 2003, the virus spread to more than 30 countries and caused disease in more than 7900 patients across five continents (2). A novel coronavirus (SCoV) was identified as the etiological agent of SARS (3, 4), and the virus causes a similar disease in cynomolgous macaques (5). Human SCoV appears to be an animal virus that crossed to humans relatively recently. Thus, identifying animals carrying the virus is of major scientific interest and public health importance. This prompted us to examine a range of domestic and wild mammals in Guangdong Province.

Because the early cases of SARS in Guangdong reportedly occurred in restaurant workers handling wild mammals as exotic food (6), our attention focused on wild animals recently captured and marketed for culinary purposes. We investigated a live-animal retail market in Shenzhen. Animals

were held, one per cage, in small wire cages. The animals sampled included seven wild, and one domestic, animal species (Table 1). They originated from different regions of southern China and had been kept in separate storehouses before arrival to the market. The animals remained in the markets for a variable period of time, and each stall holder had only a few animals of a given species. Animals from different stalls within the market were sampled. Nasal and fecal samples were collected with swabs and stored in medium 199 with bovine serum albumin and antibiotics. Where possible, blood samples were collected for serology. Before sampling, all animals were examined by a veterinary surgeon and confirmed to be free of overt disease. Serum samples were also obtained, after informed consent, from traders in animals ($n = 35$) and vegetables ($n = 20$) within the market. Sera ($n = 60$) submitted for routine laboratory tests from patients hospitalized for nonrespiratory disease in Guangdong were made anonymous and used for comparison.

Nasal and fecal swabs from 25 animals were tested for SCoV viral nucleic acid by using reverse transcription–polymerase chain reaction (RT-PCR) for the N gene of the human SCoV. Swabs from four of six Himalayan palm civets were positive in the RT-PCR assay (Table 1). All specimens were inoculated into FRhK-4 cells as previously described for virus isolation (3). A cytopathic effect was observed in cells inoculated with specimens from four Himalayan palm civets (*Paguma larvata*), two of which also positive for coronavirus in the original specimen by RT-PCR. A virus was also detected by virus isolation and direct

RT-PCR from the fecal swab of a raccoon dog (*Nyctereutes procyonoides*). No virus was detectable in six other species sampled. Electron microscopy of one infected cell supernatant (SZ16) showed viral particles with a morphology compatible with coronavirus (fig. S1). Sera from five animals had neutralizing antibody to the animal coronavirus; these were from three palm civets, a raccoon dog, and a Chinese ferret badger, respectively (Table 1).

To further validate the results from the neutralization test, a Western blot assay was used to detect SCoV-specific antibodies from these animal serum samples (Fig. 1). Indications of positive antibodies were observed from samples SZ2, SZ3, SZ11, and SZ17 (which were also positive in the neutralization assay) and from the positive control human serum. No positive signal was observed from those serum samples that were negative in the neutralization test. There was insufficient serum left over from the raccoon dog (SZ13) to be analyzed by this assay.

Table 1. Animal species tested for coronavirus detection. Abbreviations of animal species: B, beaver (*Castor fiber*); CFB, Chinese ferret-badger (*Melogale moschata*); CH, Chinese hare (*Lepus sinensis*); CM, Chinese muntjac (*Muntiacus reevesi*); DC, domestic cat (*Felis catus*); HB, hog-badger (*Arctonyx collaris*); HPC, Himalayan palm civet (*P. larvata*); RD, raccoon dog (*N. procyonoides*) (9). N, nasal sample; F, fecal sample; titer to SZ16, neutralizing antibody titer to SZ16; + denotes positive by RT-PCR or virus isolation; * denotes the PCR product or virus isolates sequenced in the study. ND, not done.

Sample number	Animal	Virus detection				Titer to SZ16
		RT-PCR		Isolation		
		N	F	N	F	
SZ1	HPC	+	*			ND
SZ2	HPC	+		+		40
SZ3	HPC	+		+	*	40
SZ4	HB					<20
SZ5	B					<20
SZ6	DC					ND
SZ7	DC					<20
SZ8	CH					ND
SZ9	CH					<20
SZ10	CM					<20
SZ11	CFB					160
SZ12	CFB					<20
SZ13	RD		+		+	≥640
SZ14	CM					<20
SZ15	B					<20
SZ16	HPC	+	+	+	+	<20
SZ17	HPC			+		≥640
SZ18	B					<20
SZ19	CH					<20
SZ20	CH					<20
SZ21	DC					<20
SZ22	DC					<20
SZ23	HB					ND
SZ24	HB					ND
SZ25	HPC			+		ND

¹Department of Microbiology, The University of Hong Kong, University Pathology Building, Queen Mary Hospital, Hong Kong Special Administrative Region (S.A.R.), of the People's Republic of China (China). ²Center for Disease Control and Prevention, Shenzhen, Guangdong Province, China. ³Department of Pathology, The University of Hong Kong, University Pathology Building, Queen Mary Hospital, Hong Kong S.A.R., China. ⁴Government Virus Unit, Department of Health, Hong Kong S.A.R., China.

*These authors contributed equally to this work.
†To whom correspondence should be addressed. E-mail: yguan@hkucc.hku.hk

Sera from humans working in the market were tested for antibody to SZ16 virus by neutralization and indirect immunofluorescence assays. Although 8 out of 20 (40%) of the wild-animal traders and 3 of 15 (20%) of those who slaughter these animals had evidence of antibody, only 1 (5%) of 20 vegetable traders was seropositive. None of these workers reported SARS-like symptoms in the past 6 months. In comparison, none of 60 control sera from patients admitted to a Guangdong

hospital for nonrespiratory diseases was seropositive (Table 2).

Two of the virus isolates (SZ3 and SZ16) isolated from the nasal swabs of palm civets were completely sequenced, and the amino acid sequence was deduced. Two other viruses were partially sequenced, from the S gene to the 3' end of the virus (GenBank accession numbers AY304486 to AY304489). Viral RNA sequences from these original swab samples from animal were confirmed in an independent laboratory (7). The full-length genome sequences had 99.8% homology to the human SCoV, which indicates that the human

and animal SCoV-like viruses were closely related. Phylogenetic analysis of the S gene of both human and animal SCoV-like viruses indicated that the animal viruses are separate from the human virus cluster (Fig. 2 and fig. S2). However, the viruses SZ1, SZ3, and SZ16 from palm civets were phylogenetically distinct. The viruses SZ3 and SZ16 had 18 nucleotide differences between them over the 29,709–base pair (bp) genome, whereas the human SCoV isolated from five geographically separate sites (GZ50, CUHK-W1, Tor-2, HKU-39848, and Urbani) differed by only 14 nucleotides (nt). Nevertheless, animal virus SZ13 (rac-




Fig. 1. Detection of antibodies against recombinant nucleocapsid protein of SCoV in animal sera by Western blot assay. Recombinant nucleocapsid protein (NP, 49.6 kDa) was used as an antigen to detect anti-SCoV antibodies in animal sera. Protein A-HRP was used as a secondary antibody, and reactive bands were visualized by the enhanced chemiluminescence Western blotting system. A serum sample from a convalescent SARS patient was used as a positive control. Blots reacted with animal (SZ2, SZ3, SZ11, SZ17, SZ7, SZ16, or SZ19) or human sera are indicated. Results from the neutralization test for SCoV-specific antibodies in these serum samples are also shown.

Fig. 2. Phylogenetic analysis of the nucleotide acid sequence of the spike gene of SCoV-like viruses. Nucleotide sequences of representative SCoV S genes (S gene coding region 21477 to 25244, 3768 bp) were analyzed. The phylogenetic tree was constructed by the neighbor-joining method with bootstrap analysis (1000 replicates) using MEGA 2 (10). Number at the nodes indicates bootstrap values in percentage. The scale bar shows genetic distance estimated using Kimura's two-parameter substitution model (11). In addition to viruses sequenced in the present study, the other sequences used in the analysis could be found in GenBank with accession number: from AY304490 to AY304495, AY278741, AY278554, AY278491, AY274119, and AY278489.




Table 2. Prevalence of antibody to animal SCoV SZ16 in humans. Controls are serum specimens from patients hospitalized for nonrespiratory diseases in Guangdong made anonymous.

Occupation	Sample numbers	Antibody positive (%)
Wild-animal trader	20	8 (40)
Slaughterer of animals	15	3 (20)
Vegetable trader	20	1 (5)
Control	60	0 (0)





Fig. 3. A 29-nt deletion in the human SCoV genome. (A) Genetic organization of SCoV-like viruses found in humans and animals. ORFs 1a and 1b, encoding the nonstructural polypeptides, and those encoding the S, E, M, and N structural proteins are indicated (green boxes). (B) Expanded view of the SCoV genomic sequence (27700 nt to 28200 nt, based on AY278554 numbering). ORFs for putative proteins and for N in human isolates are indicated as brown and green boxes, respectively (8). An extra 29-nt sequence is present downstream of the nucleotide of 27868 of the animal SCoV (based on AY278554 numbering). The presence of this 29-nt sequence in animals isolates results in fusing the ORFs 10 and 11 (top) into a new ORF (bottom; ORF10', light blue box). (C) Protein sequence alignment of ORF10 and 11 from human isolates and ORF 10' from animal isolates.

presence of this 29-nt sequence in animals isolates results in fusing the ORFs 10 and 11 (top) into a new ORF (bottom; ORF10', light blue box).

REPORTS

Table 3. Nucleotide sequence variation of the S gene of animal and human SCoV. The nucleotide residues are based on AY278554 numbering. Nonsilent mutations are highlighted in red. Dash indicates a nucleotide deletion.

Virus	Nucleotide residue																																					
	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2	2								
SZ3	C	A	T	T	C	A	T	A	T	T	C	A	G	G	G	C	A	A	G	T	G	T	T	C	C	T	C	G	T	G	C	G	C	G	C	T	G	T
SZ16	C	A	T	T	C	A	T	A	T	T	C	A	G	G	G	C	G	A	G	T	G	A	T	C	C	T	C	G	T	G	C	G	C	T	C	T	G	T
SZ1	C	A	T	T	C	A	T	A	T	T	C	A	G	G	G	C	G	A	G	T	G	A	T	C	C	T	T	G	T	G	C	G	T	T	T	T	G	T
SZ13	C	A	T	T	C	A	T	A	T	T	C	A	G	G	G	C	G	A	G	T	G	A	T	C	C	T	C	G	T	G	C	G	C	T	C	T	G	T
GZ01	C	A	T	T	C	A	C	C	T	C	C	C	A	G	G	T	G	T	C	A	G	T	T	T	T	C	C	A	C	G	T	A	C	G	C	T	A	T
GZ43	C	-	-	-	G	A	T	C	T	C	C	C	A	G	G	T	G	T	C	T	G	T	T	T	C	C	C	A	C	G	C	A	C	G	C	T	A	C
GZ60	C	-	-	-	G	A	T	C	T	C	C	C	A	G	G	T	G	T	C	T	G	T	T	T	C	C	C	A	C	G	C	A	C	G	C	T	A	C
GZ50	T	A	T	T	C	A	T	C	C	C	C	C	G	A	A	T	G	T	C	T	G	T	T	T	T	C	C	A	C	G	C	A	C	G	T	T	A	T
CUHK-W1	C	A	T	T	C	A	T	C	C	C	C	C	G	A	A	T	G	T	C	T	G	T	T	T	T	C	C	A	C	T	C	A	C	G	T	T	A	T
HKU-36871	C	A	T	T	C	A	T	C	C	C	C	C	G	A	A	T	G	T	C	T	G	T	T	T	T	C	C	A	C	T	C	A	C	G	T	T	A	T
HKU-39848	C	A	T	T	C	G	T	C	C	C	T	C	G	A	A	T	G	T	C	T	G	T	T	T	T	C	C	A	C	T	C	A	C	G	T	T	A	T
HKU-66078	C	A	T	T	C	G	T	C	C	C	T	C	G	A	A	T	G	T	C	T	G	T	T	T	T	C	C	A	C	T	C	A	C	G	T	T	A	T
HKU-65806	C	A	T	T	C	G	T	C	C	C	T	C	G	A	A	T	G	T	C	T	G	T	T	T	T	C	C	A	C	T	C	A	C	G	T	T	A	T
Urbani	C	A	T	T	C	G	T	C	C	C	T	C	G	A	A	T	G	T	C	T	G	T	T	T	T	C	C	A	C	T	C	A	C	G	T	C	A	T
Tor2	C	A	T	T	C	G	T	C	C	C	T	C	G	A	A	T	G	T	C	T	G	T	G	T	T	C	C	A	C	T	C	A	C	G	T	T	A	T

coon dog) and SZ16 (palm civet) were genetically almost identical, and transmission or contamination from one host to the other within the market cannot be excluded.

When the full genome of the animal ($n = 2$) and human ($n = 5$, see above) virus groups were compared, the most striking difference was that these human viruses have a 29-nt deletion (5'-CCTACTGGT-TACCAACCTGAATGGAATAT-3', residue 27869 to 27897) that is 246 nt upstream of the start codon of the N gene (Fig. 3). Of human SCoV sequences currently available in GenBank, there was only one (GZ01) with this additional 29-nt sequence. In addition to that, there were 43 to 57 nucleotide differences observed over the rest of the genome. Most of these differences were found in the S gene coding region. The existence of the additional 29-nt sequence in the animal viruses results in demolishing the open reading frames (ORFs) 10 and 11 (8) and merging these two ORFs into a new ORF encoding a putative protein of 122 amino acids (Fig. 3). This putative peptide has a high homology to the putative proteins encoded by ORF10 and ORF11. Because ORF11 does not have a typical transcription regulatory sequence for SCoV (8), the putative ORF11 reported by others may just be the direct result of the deletion of the 29-nt sequence. BLAST search of this peptide yields no significant match to any other known peptide. Further investigation is required to elucidate the biological significance of this finding.

When the S-gene sequences of the four animal viruses were compared with 11 human SCoV viruses, 38 nucleotide polymorphisms were noted, and 26 of them were nonsynonymous changes (Table 3). The S

genes among the four animal viruses had eight nucleotide differences, whereas there were 20 nucleotide differences among 11 human viruses. Thus, the animal viruses, although isolated from one market, are no less divergent than the human viruses isolated from Hong Kong, Guangdong, Canada, and Vietnam. However, whereas 14 (70%) of the 20 polymorphisms among the human viruses were nonsynonymous mutations, only two (25%) of the eight nucleotide substitutions within the animal viruses were. An amino acid deletion (nucleotide positions 21690 to 21692) was observed in two of the human viruses (GZ43 and GZ60). Of the 38 polymorphisms, there were 11 consistent nucleotide signatures that appeared to distinguish animal and human viruses. The observation that the human and animal viruses are phylogenetically distinct (Fig. 2) makes it highly unlikely that the SCoV-like viruses isolated in these wild animals is due to the transmission of SCoV from human to animals.

Our findings suggest that the markets provide a venue for the animal SCoV-like viruses to amplify and to be transmitted to new hosts, including humans, and this is critically important from the point of view of public health. However, it is not clear whether any one or more of these animals are the natural reservoir in the wild. It is conceivable that civets, raccoon dog, and ferret badgers were all infected from another, as yet unknown, animal source, which is in fact the true reservoir in nature. However, because of the culinary practices of southern China, these market animals may be intermediate hosts that increase the opportunity for transmission of infection to humans. Further extensive surveillance on

animals will help to better understand the animal reservoir in nature and the interspecies transmission events that led to the origin of the SARS outbreak.

References and Notes

- World Health Organization (WHO) (www.who.int/csr/sars/en/).
- WHO, Cumulative Number of Reported Probable Cases of Severe Acute Respiratory Syndrome (SARS) (www.who.int/csr/sars/country/2003_05_20/en/).
- J. S. M. Peiris *et al.*, *Lancet* **361**, 1319 (2003).
- T. G. Ksiazek *et al.*, *N. Engl. J. Med.* **348**, 1953 (2003).
- R. A. Fouchier *et al.*, *Nature* **423**, 240 (2003).
- N. S. Zhong *et al.*, *Lancet*, in press.
- K. Holmes, unpublished observations.
- M. A. Marra *et al.*, *Science* **300**, 1399 (2003).
- China species information system (www.chinabiodiversity.com).
- S. Kumara *et al.*, *Bioinformatics* **17**, 1244 (2001).
- M. Kimura, *J. Mol. Evol.* **16**, 111 (1980).
- We thank the Department of Health and Department of Agriculture of Shenzhen Government for facilitating the study. We gratefully acknowledge the encouragement and support of L. C. Tsui, Vice-Chancellor, The University of Hong Kong. We thank X. Y. Zhao from the Department of the Microbiology, The University of Hong Kong, for the excellent technical assistance. We also thank C. C. Hon and F. C. Leung from the Department of Zoology, The University of Hong Kong, and Richard Webby from St. Jude Children's Research Hospital (Memphis, TN, U.S.A.) for assistance in the phylogenetic analysis. We thank K. V. Holmes's laboratory from the Department of Microbiology, University of Colorado Health Sciences Center (Denver, CO, U.S.A.) to validate the animal viral sequences. Supported by research funding from Public Health Research (Grant A195357), the U.S. National Institute of Allergy and Infectious Diseases, the Wellcome Trust (067072/D/02/Z), and SARS research funds from The University of Hong Kong.

Supporting Online Material

www.sciencemag.org/cgi/content/full/1087139/DC1
Materials and Methods
Figs. S1 and S2
References and Notes

22 May 2003; accepted 26 August 2003

Published online 4 September 2003;

10.1126/science.1087139

Include this information when citing this paper.