

SURVIVAL AND COEXISTENCE FOR A MULTITYPE CONTACT PROCESS

J. THEODORE COX AND RINALDO B. SCHINAZI

ABSTRACT. We study the ergodic theory of a multitype contact process with equal death rates and unequal birth rates on the d -dimensional integer lattice and regular trees. We prove that for birth rates in a certain interval there is coexistence on the tree, which by a result of Neuhauser is not possible on the lattice. We also prove a complete convergence result when the larger birth rates falls outside of this interval.

1. INTRODUCTION AND RESULTS

We consider the multitype contact process on a countable set S , introduced by Neuhauser in [8]. In this paper S is either the d -dimensional integer lattice \mathbb{Z}^d , the case considered by Neuhauser, or the homogeneous connected tree \mathbb{T}_d in which each vertex has $d + 1$ neighbors, $d \geq 2$. The primary reason we are interested in the multitype contact process defined on the tree \mathbb{T}_d is that, as we show below, it exhibits phenomena that the process defined on \mathbb{Z}^d does not.

In this model each point, or site, of S is either vacant or occupied by an individual of one of two possible types. The system is described by a configuration $\xi \in \{0, 1, 2\}^S$, where $\xi(x) = 0$ means that site x is vacant, and for $i = 1, 2$, $\xi(x) = i$ means that x is occupied by an individual of type i . For $x, y \in S$ write $x \sim y$ if x, y are nearest neighbors, and define

$${}^i\xi = \{x : \xi(x) = i\}, \quad \xi \in \{0, 1, 2\}^S, \quad i = 1, 2.$$

For $x \in S$ and $\xi \in \{0, 1, 2\}^S$, let $n_i(x, \xi)$ denote the number of neighbors of x that are of type i ,

$$n_i(x, \xi) = \sum_{y \sim x} 1\{\xi(y) = i\}, \quad i = 1, 2.$$

The multitype contact process ξ_t with birth rates λ_1, λ_2 is the Feller process taking values in $\{0, 1, 2\}^S$ which makes transitions at x in configuration ξ

$$(1.1) \quad i \rightarrow 0 \text{ at rate } 1, \quad i = 1, 2,$$

$$(1.2) \quad 0 \rightarrow i \text{ at rate } \lambda_i n_i(x, \xi), \quad i = 1, 2.$$

Informally, an individual of either type dies at rate one, and for each site x and each neighboring site y , an individual of type i at x gives birth at rate λ_i to an

Date: Draft date is 06/12/2007, printed April 15, 2008.

2000 Mathematics Subject Classification. Primary: 60K35, 60G57, Secondary: 60F05, 60J80.

Key words and phrases. Contact process, trees, multitype, survival, coexistence, complete convergence.

This material is based upon work supported by the National Science Foundation under Grants No. 0505439 and No. 0701396.

individual of the same type at y provided that site is vacant. Thus the two types interact in their competition for space. Existence and uniqueness of a Feller process ξ_t determined by the above rates follows from general results of [6] (see Theorem I.3.9). We will give a “graphical construction” of ξ_t in Section 2 below.

For this process on \mathbb{Z}^d , Neuhauser proved in [8] that the two types can coexist if and only if $d \geq 3$ and $\lambda_1 = \lambda_2 > \lambda_c$ (λ_c is defined below). In [8] coexistence meant the existence of a translation invariant measure ν which is invariant for the process and concentrates on configurations with infinitely many individuals of each type. That is, letting $|\cdot|$ denote cardinality, ν should satisfy

$$(1.3) \quad \nu(\xi : |^1\xi| = |^2\xi| = \infty) = 1.$$

The non-coexistence part of this result was strengthened by Durrett and Neuhauser in [2]. By Theorem 2 there, if $\lambda_2 > \lambda_1 \vee \lambda_c$, and if the initial configuration ξ_0 has infinitely many type 2 individuals, then for every site x , $P(\xi_t(x) = 1) \rightarrow 0$ as $t \rightarrow \infty$. Consequently, there is no invariant measure ν , translation invariant or otherwise, satisfying (1.3).

Switching from \mathbb{Z}^d to \mathbb{T}_d , one can use the approach from [8] to show that if $\lambda_1 \neq \lambda_2$ then there can be no invariant measure ν which is homogeneous (invariant under the obvious shifts) and also satisfies (1.3). However, the arguments in [2], which involve a block construction, do not seem to be directly applicable on the tree, leaving open the possibility of nonhomogeneous invariant measures satisfying (1.3). We show in Theorem 1 that such measures do in fact exist provided the birth rates lie in a certain interval. In Theorem 2 we show that there is no coexistence when the birth rates are outside this interval. For this result our method of proof applies equally well on \mathbb{Z}^d .

In order to state our results it is necessary to briefly review the basic, single-type contact process ζ_t introduced by Harris in [4], treated in detail in Chapter VI of [6] and in Part I of [7]. In this process each site of S is either vacant or occupied by an individual, and ζ_t is the set of occupied sites at time t . Supposing S is either \mathbb{Z}^d or \mathbb{T}_d and $A \subset S$, the transition rates are given by

$$\begin{aligned} A &\rightarrow A \setminus \{x\} \text{ for } x \in A \text{ at rate } 1, \\ A &\rightarrow A \cup \{x\} \text{ for } x \notin A \text{ at rate } \lambda|\{y \in A : y \sim x\}|. \end{aligned}$$

For $A \subset S$ let ζ_t^A denote the process with initial state $\zeta_0^A = A$, and for $x \in S$ write ζ_t^x for $\zeta_t^{\{x\}}$. We give a construction of ζ_t in Section 2 below.

Define the two critical values $\lambda_* \leq \lambda^*$ by

$$\lambda_* = \inf\{\lambda : P(\zeta_t^x \neq \emptyset \forall t > 0) > 0\}$$

and

$$\lambda^* = \inf\{\lambda : P(x \in \zeta_t^x \text{ i.o. as } t \rightarrow \infty) > 0\}.$$

By translation invariance, the above probabilities do not depend on x . The contact process is said to *die out* if $P(|\zeta_t^x| \geq 1 \forall t > 0) = 0$, *survive weakly* if this probability is positive but $P(x \in \zeta_t^x \text{ i.o. as } t \rightarrow \infty) = 0$, and *survive strongly* if this last probability is positive. It is known that for $S = \mathbb{Z}^d$ weak survival does not occur, $\lambda_* = \lambda^*$, and the common value λ_c is finite and strictly positive. However, for the tree \mathbb{T}_d , weak survival does occur. It is known that $0 < \lambda_* < \lambda^* < \infty$, the

process dies out if $\lambda = \lambda_*$ and survives weakly if $\lambda = \lambda^*$ (see Proposition 4.39 and Theorem 4.65 in [6]). It turns out that these different possibilities for survival must be taken into account when investigating the question of coexistence for the multitype process on \mathbb{T}_d .

For the tree \mathbb{T}_d fix a site $\mathcal{O} \in \mathbb{T}_d$ and call it the root. For $x \neq y \in \mathbb{T}_d$ there is a unique sequence of distinct sites $x = x_0, x_1, \dots, x_n = y$ such that $x_{i-1} \sim x_i$ for $i = 1, \dots, n$. Let this n be the distance from x to y , $|x - y| = n$. Let B_K be the ball of radius K centered at \mathcal{O} , $B_K = \{y \in \mathbb{T}_d : |y - \mathcal{O}| \leq K\}$, and let ∂B_K denote the outer boundary, $\partial B_K = \{y \in \mathbb{T}_d : |y - \mathcal{O}| = K + 1\}$. For $x \in \mathbb{T}_d$ let $S(x)$ be the sector of the tree starting at x pointing away from the root, i.e.,

$$S(x) = \{y \in \mathbb{T}_d : |y - \mathcal{O}| = |y - x| + |x - \mathcal{O}|\}.$$

A nice set of configurations to work with is Ξ_0 , the set of configurations $\eta \in \{0, 1, 2\}^{\mathbb{T}_d}$ which satisfy:

$$(1.4) \quad \exists K < \infty \text{ such that } \eta \text{ is constant on each } S(y), y \in \partial B_K.$$

Our first results are for the case that both birth rates lie between the two contact process critical values. For this case there is coexistence, and even weak convergence starting from any configuration in Ξ_0 . For a probability measure ν on $\{0, 1, 2\}^S$, $\xi_t \Rightarrow \nu$ as $t \rightarrow \infty$ means that the finite dimensional distributions of ξ_t converge to those given by ν .

Theorem 1. *Assume $S = \mathbb{T}_d$ and $\lambda_1, \lambda_2 \in (\lambda_*, \lambda^*]$.*

(i) *If $\xi_0 \equiv i$ on $S(y)$ for some site y , then for all sites x ,*

$$(1.5) \quad \liminf_{t \rightarrow \infty} P(\xi_t(x) = i) > 0.$$

(ii) *If $\xi_0 = \eta \in \Xi_0$ then*

$$(1.6) \quad \xi_t \Rightarrow \nu_\eta \text{ as } t \rightarrow \infty$$

for some measure ν_η which is necessarily invariant for ξ_t .

(iii) *Assume $\eta \in \Xi_0$. If $|^i \eta| = \infty$, then $\nu_\eta(\xi : |^i \xi| = \infty) = 1$. If $\eta \equiv i$ on some sector $S(y)$ and $|x| \rightarrow \infty$, $x \in S(y)$, then $\nu_\eta(\xi : \xi(x) = j) \rightarrow 0$ if j is not 0 or i .*

(iv) *Assume $\eta, \eta' \in \Xi_0$. If $|\{x : \eta(x) \neq \eta'(x)\}| < \infty$ then $\nu_\eta = \nu_{\eta'}$. If $\exists y \in \mathbb{T}_d$ and $i \neq j$ such that $\eta \equiv i$ and $\eta' \equiv j$ on $S(y)$, then $\nu_\eta \neq \nu_{\eta'}$.*

Theorem 1 shows there is coexistence even for unequal birth rates provided the two rates lie in the contact process weak survival interval, and exhibits a large class of nonhomogeneous invariant measures ν_η . If $\eta \in \Xi_0$ is identically 1 on some sector and identically 2 on another, then by (ii) and (iii) above, ν_η is a nonhomogeneous invariant measure which satisfies (1.3). This means that when $\lambda_1 < \lambda_2$ and both rates lie in $(\lambda_*, \lambda^*]$, the 2's are not strong enough to drive the 1's from bounded regions of the tree, and coexistence is possible. Two-type competition models have been studied by many others, see [1], [3] and [5] for instance. The model in [5] exhibits “global” coexistence with unequal rates, in that $P(|^1 \xi_t| \geq 1, |^2 \xi_t| \geq 1 \forall t > 0) > 0$ for every initial configuration with $|^1 \xi_0| \geq 1, |^2 \xi_0| \geq 1$. Coexistence results are proved in [1] for the multitype contact process on \mathbb{Z}^d with long-range interactions, or an additional “death” mechanism. Theorem 1 may give the first

“local” coexistence result for a nearest-neighbor interaction model with equal death rates and unequal birth rates.

It is a different story when $\lambda_1 \neq \lambda_2$ with one or both rates larger than λ^* . In this case coexistence, even in the weak sense of (1.7) below, is not possible.

Theorem 2. *Assume S is \mathbb{Z}^d or \mathbb{T}_d and $\lambda_2 > \lambda_1 \vee \lambda^*$. For any initial configuration ξ_0 and $x \in S$,*

$$(1.7) \quad \lim_{t \rightarrow \infty} P(\xi_t(x) = 1 \text{ and } |{}^2\xi_t| \geq 1) = 0.$$

Consequently, $\lim_{t \rightarrow \infty} P(\xi_t(x) = 1) = 0$ if $|{}^2\xi_0| = \infty$.

For $S = \mathbb{Z}^d$, (1.7) follows from the results in Section 3 of [2]. We include this case in the statement of Theorem 2 since our proof for $S = \mathbb{T}_d$ applies equally well to the lattice case, and is simpler than the one in [2]. The lack of coexistence in (1.7) means that an invariant measures can concentrate on only one type, and allows us to prove that the process converges weakly from any initial configuration. We need some additional notation and information about the basic contact process before stating our results.

For the single-type contact process ζ_t with birth rate λ define the survival probabilities $\alpha_A = \alpha_A(\lambda)$ by

$$(1.8) \quad \alpha_A = P(\zeta_t^A \neq \emptyset \forall t \geq 0), \quad A \subset S,$$

and let $\alpha = \alpha_{\{x\}}$, which by translation invariance does not depend on x .

It is well known, (see (I.1.4) of [6]), that there is a probability measure $\bar{\nu} = \bar{\nu}_\lambda$ called the upper invariant measure such that

$$(1.9) \quad \zeta_t^S \Rightarrow \bar{\nu}_\lambda \text{ as } t \rightarrow \infty.$$

Letting δ_\emptyset be the unit point mass on the empty set, $\bar{\nu} \neq \delta_\emptyset$ if and only if $\lambda > \lambda_*$. The *complete convergence theorem* for ζ_t is the statement

$$(1.10) \quad \zeta_t^A \Rightarrow (1 - \alpha_A)\delta_\emptyset + \alpha_A\bar{\nu} \text{ as } t \rightarrow \infty \quad \forall A \subset S.$$

It is known that (1.10) holds for both \mathbb{Z}^d and \mathbb{T}_d if $\lambda > \lambda^*$ (see Theorems 1.2.27 and 1.4.70 in [6]).

We show here that there is an analogous theorem for the multitype contact process if $\lambda_2 > \lambda_1 > \lambda^*$. To state it we must define appropriate survival probabilities and “upper invariant measures.” For birth rates λ_1, λ_2 and configurations $\eta \in \{0, 1, 2\}^S$ let $\xi_0 = \eta$ and define $\alpha_\eta^i = \alpha_\eta^i(\lambda_1, \lambda_2)$, $i = 1, 2$, by

$$(1.11) \quad \begin{aligned} \alpha_\eta^1 &= P({}^1\xi_t \neq \emptyset \forall t \geq 0 \text{ and } {}^2\xi_t = \emptyset \text{ eventually}), \\ \alpha_\eta^2 &= P({}^2\xi_t \neq \emptyset \forall t \geq 0). \end{aligned}$$

We need probability measures $\bar{\nu}^i = \bar{\nu}_{\lambda_i}^i$ on $\{0, 1, 2\}^S$ which correspond to $\bar{\nu}_{\lambda_i}$ and concentrate on configurations in which all individuals are of type i . These measures are defined by the requirements

$$\bar{\nu}^1(\xi : {}^2\xi = \emptyset) = \bar{\nu}^2(\xi : {}^1\xi(x) = \emptyset) = 1$$

and

$$\bar{\nu}^i(\xi : {}^i\xi \cap A \neq \emptyset) = \bar{\nu}_{\lambda_i}(\zeta : \zeta \cap A \neq \emptyset), \quad A \subset S, i = 1, 2.$$

With these definitions in place we can now state our complete convergence theorem for ξ_t when $\lambda_2 > \lambda_1 > \lambda^*$. In the following let δ_0 denote the measure on $\{0, 1, 2\}^S$ which concentrates on the single configuration $\xi \equiv 0$.

Theorem 3. *Assume S is \mathbb{Z}^d or T^d , $\lambda_2 > \lambda_1 > \lambda^*$ and $\xi_0 = \eta$. Then*

$$(1.12) \quad \xi_t \Rightarrow (1 - \alpha_\eta^1 - \alpha_\eta^2)\delta_0 + \alpha_\eta^1\bar{\nu}^1 + \alpha_\eta^2\bar{\nu}^2 \quad \text{as } t \rightarrow \infty.$$

Given Theorem 2, this result is not surprising. If the 2's survive then the 1's are driven out of bounded regions, so the 2's in effect form a single type contact process and (1.10) takes over. If there are finitely many 2's which die out while the 1's manage to survive, then the 1's form a single type contact process and again (1.10) takes over.

The complete convergence theorem for the contact process (1.10) does not hold on \mathbb{T}_d for $\lambda \in (\lambda_*, \lambda^*]$. In this case the contact process has a wide variety of invariant measures (see Theorems I.4.107 and I.4.121 of [6]) and hence possible limits for ζ_t . We cannot expect (1.12) to hold as stated if $\lambda_1 \leq \lambda^* < \lambda_2$. However, if we restrict ξ_0 to configurations for which the corresponding contact process of 1's converges weakly, then ξ_t also converges weakly.

Theorem 4. *Assume $S = T^d$, $\lambda_2 > \lambda^* \geq \lambda_1$ and $\xi_0 = \eta$. Let ζ_t be the single-type contact process with birth rate λ_1 and initial state $\zeta_0 = {}^1\eta$, and assume that $\zeta_t \Rightarrow \mu$ as $t \rightarrow \infty$. Then*

$$(1.13) \quad \xi_t \Rightarrow (1 - \alpha_\eta^2)\bar{\mu}^1 + \alpha_\eta^2\bar{\nu}^2 \quad \text{as } t \rightarrow \infty,$$

where $\mu^1(\xi : {}^2\xi = \emptyset) = 1$ and $\mu^1(\xi : {}^1\xi \cap A \neq \emptyset) = \mu(\zeta : \zeta \cap A \neq \emptyset)$.

For $S = \mathbb{Z}^d$, the conclusions of Theorems 2 and 3 can be derived from results in [2] (Lemma 3 and the construction in Section 3 there). However, our methods are simpler and should apply without much change to other choices for S , including some periodic graphs.

As previously noted, for $S = \mathbb{Z}^d$ and $\lambda_1 = \lambda_2 > \lambda_c$, it was shown in [8] that there is coexistence for $d \geq 3$ but not for $d \leq 2$. Presumably the arguments for these results can be adapted to handle the tree \mathbb{T}_d , where one expects there should be coexistence for $\lambda_1 = \lambda_2 > \lambda^*$.

In the next section we construct our processes using the standard “graphical construction” via Poisson processes. The construction naturally contains various couplings and dual processes used in our proofs. In Sections 3-6 we prove Theorems 1-4, respectively.

We note that our main tool is the *ancestor duality* introduced by Neuhauser in [8].

2. CONSTRUCTION AND DUALITY

We start by constructing our process using Harris' graphical method, assuming from now on that

$$(2.1) \quad \lambda_1 \leq \lambda_2.$$

The construction takes place in the space-time set $S \times [0, \infty)$ using independent families of Poisson processes. For $x \in S$ let $\{T_n^x : n \geq 1\}$ be the arrival times of a

Poisson process with rate 1. At the times T_n^x we put a δ at site x to indicate that there is a death at x : if site x is occupied by either type it becomes vacant at that time. For all pairs of nearest neighbors $x, y \in S$ let $\{B_n^{x,y} : n \geq 1\}$ be the arrival times of a Poisson process with rate λ_2 . At the times $B_n^{x,y}$ we do two things. We draw an arrow from site x to site y , and with probability $1 - \lambda_1/\lambda_2$, independently of everything else, label the arrow with a “2” (and otherwise do not label the arrow). If there is a 2 at x and y is vacant at that time then there is a birth of a 2 at y . If there is a 1 at x and y is vacant we put a 1 at y provided the arrow does not have a 2 on it. Thus, the 2-arrows are really “2-only” arrows. If $\lambda_1 = \lambda_2$ then no arrow is marked with a 2. The poisson processes $T^x, B^{x,y}$, $x, y \in S$, are all independent of one another.

For sites $x, y \in S$ and times $0 < s \leq t$, we say there is a path up from (x, s) to (y, t) if there is a sequence of times $t_0 = s < t_1 < t_2 < \dots < t_n = t$ and a sequence of sites $x_0 = x, x_1, \dots, x_n = y$ such that for $i = 1, 2, \dots, n$, $x_{i-1} \sim x_i$, there is an arrow from x_{i-1} to x_i at time t_i and the time segments $\{x_i\} \times (t_{i-1}, t_i)$ do not contain any δ 's. By default there is always a path up from (x, t) to (x, t) . A path up which has at least one arrow labeled 2 will be called a “2-path,” and a path with no arrows labeled 2 will be called a “1-path.” Note that 1's propagate *only* along 1-paths, but 2's propagate along both 1-paths and 2-paths. For $s < t$ there is an i -path down from (y, t) to (x, s) if and only if there is an i -path up from (x, s) to (y, t) . Given an initial configuration ξ_0 we may construct $\xi_t, t \geq 0$ from our Poisson processes by following paths from occupied sites forward in time, using δ 's for deaths and arrows for births, as appropriate.

We now define several “reverse” time dual processes, starting with the simplest. For $x \in S, t > 0, i = 1, 2$ and $0 \leq s \leq t$, define

$$D_s^{i,(x,t)} = \{y \in S : \text{there is an } i\text{-path down from } (x, t) \text{ to } (y, t - s)\}$$

and $D_s^{(x,t)} = D_s^{1,(x,t)} \cup D_s^{2,(x,t)}$. The *ancestor process* introduced in [8] is more complicated. Fix $x \in S$ and $t > 0$, and consider $D_s^{(x,t)}, 0 \leq s \leq t$. If $D_s^{(x,t)}$ is not empty then the sites in $D_s^{(x,t)}$ are the possible *ancestors* at (forward) time $t - s$ of (x, t) , which can be arranged in decreasing order of priority, $(a_1(s), a_2(s), \dots, a_n(s))$ for some n , with $a_1(s)$ denoting the *primary* ancestor (see [8]). The j th ancestor a_j is associated with a path up from $(a_j, t - s)$ to (x, t) , which may or may not contain an arrow labeled 2, blocking propagation of 1's. We will use an equivalent but slightly different formulation of this process, which we now describe.

An ancestor configuration $\hat{\xi}$ is either the empty set, or a sequence of pairs $((a_1, b_1), \dots, (a_n, b_n))$ for some $n \geq 1$, where each $a_j \in S$ and b_j is either 1 or 2. The ancestor process $\hat{\xi}_s^{(x,t)}, 0 \leq s \leq t$ is a Markov process defined as follows. First, put $\hat{\xi}_0^{(x,t)} = ((x, 1))$. Suppose now that $s < t$ and $\hat{\xi}_u^{(x,t)}$ has been defined for $u \in [0, s]$. If $\hat{\xi}_s^{(x,t)} = \emptyset$ put $\hat{\xi}_v^{(x,t)} = \emptyset$ for all $s < v \leq t$. If $\hat{\xi}_s^{(x,t)} = ((a_1, b_1), \dots, (a_n, b_n))$ for some $n \geq 1$, let u be the the smallest time larger than s at which an event occurs at (forward) time $t - u$ affecting any of the a_j . If there is no such $u \leq t$ put $\hat{\xi}_v^{(x,t)} = \hat{\xi}_s^{(x,t)}$ for all $s < v \leq t$, we are done. Now suppose $u < t$.

- (1) If the event affecting a_j at time $t - u$ is an arrow pointing from some site a to a_j , insert (a, b) into $((a_1, b_1), \dots, (a_n, b_n))$ after each (a_i, b_i) such that $a_i = a_j$.
 - If the arrow from a to a_j is labeled 2, set $b = 2$ for each of these insertions.
 - If the arrow is unlabeled, set $b = 1$, except for insertions after any (a_i, b_i) with $b_i = 2$, in which case set $b = 2$.
- (2) If the event affecting a_j is a δ , then delete each (a_i, b_i) from $((a_1, b_1), \dots, (a_n, b_n))$ with $a_i = a_j$.

Let $\widehat{\xi}_u^{(x,t)}$ be the resulting sequence, setting $\widehat{\xi}_u^{(x,t)} = \emptyset$ if all the a_i were deleted. Iteration of this procedure defines $\widehat{\xi}_v^{(x,t)}$ for all $v \in [0, t]$. For an ancestor configuration $\widehat{\xi} = ((a_1, b_1), \dots, (a_n, b_n))$ let $\text{supp}(\widehat{\xi}) = \{a_j : 1 \leq j \leq n\}$, so that $\text{supp}(\widehat{\xi}_s^{(x,t)}) = D_s^{(x,t)}$. We say that the j th ancestor a_j is *1-blocked* if $b_j = 2$. If $\lambda_1 = \lambda_2$ then there are no 1-blocked ancestors and the b_j can be dispensed with.

The duality equation relating $\widehat{\xi}_s^{(x,t)}$ and ξ_{t-s} is

$$(2.2) \quad \xi_t(x) = \Psi(x, \widehat{\xi}_s^{(x,t)}, \xi_{t-s}), \quad s \in [0, t],$$

where $\Psi(x, \widehat{\xi}, \xi)$ is the function of sites $x \in S$, ancestor configurations $\widehat{\xi}$, and configurations $\xi \in \{0, 1, 2\}^S$ defined as follows. If $\widehat{\xi} = \emptyset$, put $\Psi(x, \widehat{\xi}, \xi) = 0$. Otherwise, $\widehat{\xi} = ((a_1, b_1), \dots, (a_n, b_n))$ for some $n \geq 1$, and we start checking the ancestors one at a time. If $\xi(a_1) = 2$ set $\Psi(x, \widehat{\xi}, \xi) = 2$ indicating a 2 propagates up. If $\xi(a_1) = 1$ and $b_1 = 1$, set $\Psi(x, \widehat{\xi}, \xi) = 1$ indicating a 1 propagates up. Now suppose either $\xi(a_1) = 1$ and $b_1 = 2$, or $\xi(a_1) = 0$. If $n = 1$ set $\Psi(x, \widehat{\xi}, \xi) = 0$. Otherwise $n \geq 2$, and we consider (a_2, b_2) and proceed as with (a_1, b_1) , either setting $\Psi(x, \widehat{\xi}, \xi)$ equal to 1 or 2 or exhausting the set of ancestors completely, in which case we set $\Psi(x, \widehat{\xi}, \xi) = 0$. The duality equation (2.2) holds because it holds at time $s = 0$, and each transition preserves its validity. For an example, see Figure 1, in which the solid circles indicate deaths, $\xi_0(a) = 2$, $\xi_0(b) = 1$, $\xi_0(c) = 0$, $\xi_0(d) = 1$, $\xi_0(e) = 2$, $\widehat{\xi}_s^{(c,t)} = ((b, 1), (a, 2), (c, 1), (d, 1), (e, 2), (d, 2))$, $\xi_{t-s}(a) = \xi_{t-s}(c) = \xi_{t-s}(e) = 1$, $\xi_{t-s}(b) = \xi_{t-s}(d) = 0$, and $\Psi(c, \widehat{\xi}_s^{(c,t)}, \xi_{t-s}) = 1$.

The reverse time ancestor processes $\widehat{\xi}_s^{(x,t)}$ are defined only for bounded time intervals. However, as in [8], we can switch to forward time, and define ancestor processes $\bar{\xi}_s^{(x,t)}$, $s \geq t$ in the analogous way, so that the laws of $\bar{\xi}_s^{(x,0)}$, $s \in [0, t]$ and $\widehat{\xi}_s^{(x,t)}$, $s \in [0, t]$ are the same. Thus, in Figure 1, $\bar{\xi}_t^{(d,t-s)} = ((d, 1), (e, 2))$. We will write $\bar{\xi}_s^x$ for $\bar{\xi}_s^{(x,0)}$.

Our construction also contains various couplings with the single-type contact process. Consider $x \in S$ and $t > 0$. For $s \geq t$ let $\zeta_s^{1,(x,t)}$ be the set of sites y such that there is a 1-path up from (x, t) to (y, s) , and let $\zeta_s^{2,(x,t)}$ be the set of sites y such that there is either a 1-path or 2-path up from (x, t) to (y, s) . Write $\zeta_s^{i,x}$ for $\zeta_s^{i,(x,0)}$ and $\zeta_s^{i,A}$ for $\cup_{x \in A} \zeta_s^{i,x}$, $i = 1, 2$. Each process $\zeta_s^{i,x}$, $s \geq 0$ is a single-type contact processes with birth rate λ_i . Also, $D_s^{1,(x,t)}$, $0 \leq s \leq t$ has the same law as $\zeta_s^{1,x}$, $0 \leq s \leq t$, and $D_s^{(x,t)}$, $0 \leq s \leq t$ has the same law as $\zeta_s^{2,x}$, $0 \leq s \leq t$. Finally, we make the observation that $\text{supp}(\bar{\xi}_s^{(x,t)})$, $s \geq t$ and $\zeta_s^{2,(x,t)}$, $s \geq t$ have the same law.




FIGURE 1.

A word on notation. Throughout we will use ξ, η for elements of $\{0, 1, 2\}^S$, ζ for subsets of S , and $\bar{\xi}$ and $\hat{\xi}$ for ancestor configurations.

3. PROOF OF THEOREM 1

Recall that (2.1) is in force.

Proof of (1.5). It suffices to assume that $\xi \equiv 1$ on $S(y)$ for some neighbor y of \mathcal{O} , and show that there is some $x \in S(y)$ such that

$$(3.1) \quad \liminf_{t \rightarrow \infty} P(\xi_t(x) = 1) > 0.$$

We start with a simple consequence of the duality equation (2.2): for any $x \in S$, then

$$(3.2) \quad \{\emptyset \neq D_t^{1,(x,t)} \text{ and } D_t^{(x,t)} \subset S(y)\} \subset \{\xi_t(x) = 1\}.$$

Rinaldo: delete explanation?

This follows because on the event on the left side $D_t^{(x,t)} = \text{supp}(\hat{\xi}_t^{(x,t)}) = ((a_1, b_1), \dots, (a_n, b_n))$ for some $n \geq 1$, $\xi_0(a_i) = 1$ for each i , and since $D_t^{1,(x,t)}$ is nonempty, at least one of the a_i is not 1-blocked. Therefore, $\Psi(x, \hat{\xi}_t^{(x,t)}, \xi_0) = 1$, so $\xi_t(x) = 1$ by (2.2).

By (3.2) we have

$$\begin{aligned} P(\xi_t(x) = 1) &\geq P(D_t^{1,(x,t)} \neq \emptyset) - P(D_t^{(x,t)} \not\subset S(y)) \\ &= P(\zeta_t^{1,x} \neq \emptyset) - P(\zeta_t^{2,x} \not\subset S(y)) \\ &\geq P(\zeta_s^{1,x} \neq \emptyset \text{ for all } s \geq 0) - P(\zeta_s^{2,x} \cap S^c(y) \neq \emptyset \text{ for some } s \geq 0). \end{aligned}$$

Since $\lambda_1 > \lambda_*$, the survival probability $\alpha(\lambda_1) = \alpha_{\{x\}}(\lambda_1)$ (recall (1.8)) is positive. For $x \in S(y)$, the fact that 2's spread only by nearest neighbor contact implies

$$\{\zeta_s^{2,x} \cap S^c(y) \neq \emptyset \text{ for some } s \geq 0\} = \{\mathcal{O} \in \zeta_s^{2,x} \text{ for some } s \geq 0\}.$$

We can now make use of Theorem I.4.65 of [6], which states that if $\lambda_2 \leq \lambda^*$ then

$$(3.3) \quad P(\mathcal{O} \in \zeta_s^{2,x} \text{ for some } s \geq 0) \leq (1/\sqrt{d})^{|x-\mathcal{O}|}.$$

(All that is important for us about this bound is that it tends to 0 as $|x-\mathcal{O}| \rightarrow \infty$.)

By combining the above it follows that for all $x \in S(y)$ and $t \geq 0$,

$$(3.4) \quad P(\xi_t(x) = 1) \geq \alpha(\lambda_1) - (1/\sqrt{d})^{|x-\mathcal{O}|},$$

which is certainly positive for $x \in S(y)$ with $|x-\mathcal{O}|$ sufficiently large. \square

Before continuing with the proof of Theorem 1 we prove a lemma which shows that limits of certain finite dimensional distributions for ξ_t exist when the dual lands in a region where the initial state is constant. For $x \in \mathbb{T}_d$ and $t > 0$ let $\tau_K^{(x,t)} = \inf\{s \in [0, t] : D_s^{(x,t)} \cap B_K \neq \emptyset\}$, with $\inf(\emptyset) = \infty$ (so $\tau_K^{(x,t)} = \infty$ simply means $D_s^{(x,t)} \cap B_K = \emptyset \forall s \leq t$). Switching to forward time, let $\tau_K^{i,x} = \inf\{s \geq 0 : \zeta_s^{i,x} \cap B_K \neq \emptyset\}$ and define the lifetimes $\sigma^{i,x} = \inf\{s \geq 0 : \zeta_s^{i,x} = \emptyset\}$, $i = 1, 2$.

Lemma 1. *Assume that $y \in \partial B_K$, ξ_0 is constant on $S(y)$, and A_0, A_1, A_2 are finite disjoint subsets of $S(y)$. Then as $t \rightarrow \infty$,*

$$(3.5) \quad P(\tau_K^{(z,t)} = \infty \text{ and } \xi_t(z) = j \forall z \in A_j, j = 0, 1, 2) \rightarrow \phi(A_0, A_1, A_2)$$

for some $\phi(A_0, A_1, A_2)$.

Proof. Let $A = A_0 \cup A_1 \cup A_2 \subset S(y)$. (i) Suppose $\xi_0 \equiv 0$ on $S(y)$. For $z \in A \subset S(y)$, if $\tau_K^{(z,t)} = \infty$, then $\text{supp}(\widehat{\xi}_t^{(z,t)}) \subset S(y)$. Consequently, no ancestor can land on a site occupied by a 1 or 2, implying $\xi_t(z)$ cannot be 1 or 2. This means the left side of (3.5) is zero if either A_1 or A_2 is nonempty. If $A_1 = A_2 = \emptyset$, the left side of (3.5) equals

$$P(\tau_K^{(z,t)} = \infty \forall z \in A) = P(\tau_K^{2,z} > t \forall z \in A) \rightarrow P(\tau_K^{2,z} = \infty \forall z \in A)$$

as $t \rightarrow \infty$.

(ii) Suppose $\xi_0 \equiv 2$ on $S(y)$. Again, $\tau_K^{(z,t)} = \infty$ implies $\text{supp}(\widehat{\xi}_t^{(z,t)}) \subset S(y)$, so now the left side of (3.5) is zero unless $A_1 = \emptyset$. In this case, $\xi_t(z) = 2$ if and only if $D_t^{(z,t)} \neq \emptyset$. Therefore, the left side of (3.5), for $A_1 = \emptyset$, equals

$$\begin{aligned} P(\tau_K^{(z,t)} = \infty \forall z \in A, D_t^{(z,t)} = \emptyset \forall z \in A_0, D_t^{(z,t)} \neq \emptyset \forall z \in A_2) \\ = P(\tau_K^{2,z} > t \forall z \in A, \sigma^{2,z} \leq t \forall z \in A_0, \sigma^{2,z} > t \forall z \in A_2) \\ \rightarrow P(\tau_K^{2,z} = \infty \forall z \in A, \sigma^{2,z} < \infty \forall z \in A_0, \sigma^{2,z} = \infty \forall z \in A_2) \end{aligned}$$

as $t \rightarrow \infty$.

(iii) Suppose $\xi_0 \equiv 1$ on $S(y)$. Now the left side of (3.5) is zero unless $A_2 = \emptyset$, and in this case $\xi_t(z) = 1$ if and only if $D_t^{1,(z,t)} \neq \emptyset$. Thus, if $A_2 = \emptyset$, the left side

of (3.5) equals

$$\begin{aligned} P(\tau_K^{(z,t)} = \infty \forall z \in A, D_t^{1,(z,t)} = \emptyset \forall z \in A_0, D_t^{1,(z,t)} \neq \emptyset \forall z \in A_1) \\ = P(\tau_K^{2,z} > t \forall z \in A, \sigma^{1,z} \leq t \forall z \in A_0, \sigma^{1,z} > t \forall z \in A_1) \\ \rightarrow P(\tau_K^{2,z} = \infty \forall z \in A, \sigma^{1,z} < \infty \forall z \in A_0, \sigma^{1,z} = \infty \forall z \in A_1) \end{aligned}$$

as $t \rightarrow \infty$. This proves the lemma. \square

Proof of Theorem 1(ii). Assume $\xi_0 \in \Xi_0$, and K is such that ξ_0 is constant on each $S(y), y \in \partial B_K$. We will prove that for any $x \in \mathbb{T}_d, a \in \{0, 1, 2\}$,

$$(3.6) \quad \lim_{t \rightarrow \infty} P(\xi_t(x) = a) \text{ exists.}$$

Afterwards we will show how to modify the proof to handle convergence of all finite dimensional distributions.

Let $L^{(x,t)}$ be the last time up to time t the dual $D_s^{(x,t)}$ starting from (x,t) contains some point of B_K , $L^{(x,t)} = \sup\{s \leq t : B_K \cap D_s^{(x,t)} \neq \emptyset\}$, with $\sup(\emptyset) = 0$. We will prove (3.6) using a decomposition based on the value of $L^{(x,t)}$. The case $L^{(x,t)} = 0$ is easily handled. If $x \in B_K$ then necessarily $L^{(x,t)} > 0$, while for $x \notin B_K$, $\{L^{(x,t)} = 0\} = \{\tau_K^{(x,t)} = \infty\}$, and so Lemma 1 implies

$$(3.7) \quad \lim_{t \rightarrow \infty} P(\xi_t(x) = a, L^{(x,t)} = 0) \text{ exists.}$$

Now suppose $L^{(x,t)} = s$ for some $s \in (0, t)$, which occurs when:

- (1) $\widehat{\xi}_{s-}^{(x,t)} = \widehat{\xi}'$ for some $\widehat{\xi}'$ containing a single $w \in B_K$ as one or more of the ancestors of (x,t) ,
- (2) there is a δ at this site w at (forward) time $t - s$,
- (3) $\widehat{\xi}_s^{(x,t)} = \widehat{\xi}$ where $\widehat{\xi}$ is obtained from $\widehat{\xi}'$ by removing each (a_i, b_i) with $a_i = w$,
- (4) $D_u^{(z,t-s)} \cap B_K = \emptyset$ for all $z \in \text{supp}(\widehat{\xi})$ and $u \in [0, t - s]$.

Using the convention above for $\widehat{\xi}'$ and $\widehat{\xi}$, the duality equation (2.2) implies that if $0 < t_0 < t$ we have

$$\begin{aligned} P(\xi_t(x) = a, 0 < L^{(x,t)} < t_0) &= \int_0^{t_0} P(L^{(x,t)} \in ds, \xi_t(x) = a) \\ &= \sum_{\widehat{\xi}'}^{(K)} \int_0^{t_0} \int P(L^{(x,t)} \in ds, \widehat{\xi}_{s-}^{(x,t)} = \widehat{\xi}', \xi_{t-s} \in d\xi) 1\{\Psi(x, \widehat{\xi}, \xi) = a\}, \end{aligned}$$

where the sum $\sum^{(K)}$ is over $\widehat{\xi}'$ such that $\text{supp}(\widehat{\xi}') \cap B_K$ is a single site. By independence of disjoint space-time regions of the Poisson processes used in the construction in Section 2, and the fact that δ 's occur at rate one, the above equals

$$\begin{aligned} \sum_{\widehat{\xi}'}^{(K)} \int_0^{t_0} \int P(\widehat{\xi}_s^{(x,t)} = \widehat{\xi}') \\ P(\tau_K^{(z,t-s)} = \infty \forall z \in \text{supp}(\widehat{\xi}), \xi_{t-s} \in d\xi) 1\{\Psi(x, \widehat{\xi}, \xi) = a\} ds. \end{aligned}$$

For fixed x and $\widehat{\xi}$, $\Psi(x, \widehat{\xi}, \xi)$ depends on ξ only through the values $\xi(z), z \in \text{supp}(\widehat{\xi})$. Hence we can define $\Pi(x, a, \widehat{\xi})$ to be the set of partitions $\mathcal{A} = (A_0, A_1, A_2)$ of $\text{supp}(\widehat{\xi})$ such that for all $\xi, \Psi(x, \widehat{\xi}, \xi) = a$ if and only if $\xi \equiv j$ on $A_j, j = 0, 1, 2$

for some $\mathcal{A} \in \Pi(x, a, \widehat{\xi})$. Therefore, summing over the possible values of ξ_{t-s} on $\text{supp}(\widehat{\xi})$, and replacing $P(\widehat{\xi}_s^{(x,t)} = \widehat{\xi}')$ with $P(\bar{\xi}_s^x = \widehat{\xi}')$, we have

$$(3.8) \quad P(\xi_t(x) = a, 0 < L^{(x,t)} \leq t_0) = \sum_{\widehat{\xi}'}^{(K)} \sum_{\mathcal{A} \in \Pi(x, a, \widehat{\xi})} \int_0^{t_0} P(\bar{\xi}_s^x = \widehat{\xi}') q_t(s, \mathcal{A}) ds$$

where

$$(3.9) \quad q_t(s, \mathcal{A}) = P\left(\tau_K^{(z, t-s)} = \infty \text{ and } \xi_{t-s}(z) = j \forall z \in A_j, j = 0, 1, 2\right).$$

It is time to use the fact that $\xi_0 \in \Xi_0$. Let S^i be the union of all $S(y)$, $y \in \partial B_K$ such that $\xi_0 \equiv i$ on $S(y)$, and for the sets A_j above let $A_j^i = A_j \cap S^i$. Since the S^i are disjoint, and the duals $D_u^{(z, t-s)}$ do not leave their respective sectors S^i on $\{\tau_K^{(z, t-s)} = \infty\}$, independence of disjoint space-time regions implies

$$\begin{aligned} q_t(s, \mathcal{A}) &= P\left(\prod_{i=0}^2 \{\tau_K^{(z, t-s)} = \infty \text{ and } \xi_{t-s}(z) = j \forall z \in A_j^i, j = 0, 1, 2\}\right) \\ &= \prod_{i=0}^2 P(\tau_K^{(z, t-s)} = \infty \text{ and } \xi_{t-s}(z) = j \forall z \in A_j^i, j = 0, 1, 2). \end{aligned}$$

By Lemma 1, the above product converges for fixed s as $t \rightarrow \infty$ to some $\phi(\mathcal{A}) = \prod_{i=0}^2 \phi(A_0^i, A_1^i, A_2^i)$. Since each $\Pi(x, a, \widehat{\xi}')$ is finite, this implies

$$(3.10) \quad \lim_{t \rightarrow \infty} \sum_{\mathcal{A} \in \Pi(x, a, \widehat{\xi})} q_t(s, \mathcal{A}) = \sum_{\mathcal{A} \in \Pi(x, a, \widehat{\xi})} \phi(\mathcal{A})$$

for fixed $s, \widehat{\xi}'$.

Since $\sum_{\mathcal{A} \in \Pi(x, a, \widehat{\xi})} q_t(s, \mathcal{A}) \leq 1$ and $\sum_{\widehat{\xi}'}^{(K)} \int_0^{t_0} P(\bar{\xi}_s^x = \widehat{\xi}') ds \leq t_0$, the dominated convergence theorem can be applied in (3.8), so that (3.10) implies

$$(3.11) \quad \lim_{t \rightarrow \infty} P(\xi_t(x) = a, 0 < L^{(x,t)} \leq t_0) = \sum_{\widehat{\xi}'}^{(K)} \sum_{\mathcal{A} \in \Pi(x, a, \widehat{\xi})} \phi(\mathcal{A}) \int_0^{t_0} P(\bar{\xi}_s^x = \widehat{\xi}') ds.$$

The contribution to (3.6) of $L^{(x,t)} > t_0$ for large t_0 is negligible, since $\lambda_2 \leq \lambda^*$ implies

$$(3.12) \quad \limsup_{t_0 \rightarrow \infty} \sup_{t \geq t_0} P(t_0 < L^{(x,t)} \leq t) \leq \limsup_{t_0 \rightarrow \infty} P(\zeta_s^{2,x} \cap B_K \neq \emptyset \text{ for some } s \geq t_0) = 0.$$

We have thus proved

$$\lim_{t \rightarrow \infty} P(\xi_t(x) = a, 0 < L^{(x,t)} \leq t) = \sum_{\widehat{\xi}'}^{(K)} \sum_{\mathcal{A} \in \Pi(x, a, \widehat{\xi})} \phi(\mathcal{A}) \int_0^\infty P(\bar{\xi}_s^x = \widehat{\xi}') ds,$$

and in view of (3.7) this implies (3.6) must hold.

More generally, let $\Gamma = \{x_1, \dots, x_m\} \subset \mathbb{T}_d$, and $a : \Gamma \rightarrow \{0, 1, 2\}$. We claim that

$$(3.13) \quad \lim_{t \rightarrow \infty} P(\xi_t(x) = a(x) \forall x \in \Gamma) \text{ exists.}$$

First, let $L^{\Gamma,t} = \max\{L^{(x,t)}, x \in \Gamma\}$, and use Lemma 1 to obtain

$$\lim_{t \rightarrow \infty} P(\xi_t(x) = a(x) \forall x \in \Gamma, L^{\Gamma,t} = 0) \text{ exists.}$$

Next, consider $P(\xi_t(x) = a(x) \forall x \in \Gamma, 0 < L^{\Gamma,t} \leq t_0)$, and decompose this event according to the value of $L^{\Gamma,t}$,

$$\begin{aligned} & P(\xi_t(x) = a(x) \forall x \in \Gamma, 0 < L^{\Gamma,t} \leq t_0) \\ &= \sum_{\widehat{\xi}_1, \dots, \widehat{\xi}_m}^{(K)} \int_0^{t_0} \int P(L^{\Gamma,t} \in ds, \widehat{\xi}_{s-}^{(x_i,t)} = \widehat{\xi}_i, 1 \leq i \leq m, \xi_{t-s} \in d\xi) \times \\ & \quad \prod_{i=1}^m 1\{\Psi(x_i, \widehat{\xi}_i, \xi) = a(x_i)\} \end{aligned}$$

where the sum $\sum^{(K)}$ is over $\widehat{\xi}_1, \dots, \widehat{\xi}_m$ such that $B_K \cap (\cup_i \text{supp}(\widehat{\xi}_i))$ is a single site $w \in B_K$. As before, for each i , $\widehat{\xi}_i$ is obtained from $\widehat{\xi}_i^t$ by removing all (a, b) with $a = w$, if any. By independence of disjoint space-time regions, the above equals

$$\begin{aligned} & \sum_{\widehat{\xi}_1, \dots, \widehat{\xi}_m}^{(K)} \int_0^{t_0} \int P(\widehat{\xi}_s^{(x_i,t)} = \widehat{\xi}_i, 1 \leq i \leq m) \times \\ & P(\tau_K^{(z,t-s)} = \infty \forall z \in \text{supp}(\cup_i \widehat{\xi}_i), \xi_{t-s} \in d\xi) \prod_{i=1}^m 1\{\Psi(x_i, \widehat{\xi}_i, \xi) = a(x_i)\} ds. \end{aligned}$$

Let $\Pi(\Gamma, a, \widehat{\xi}_1, \dots, \widehat{\xi}_m)$ be the set of partitions $\mathcal{A} = (A_0, A_1, A_2)$ of $\cup_i \text{supp}(\widehat{\xi}_i)$ such that $\Psi(x_i, \widehat{\xi}_i, \xi) = a(x_i) \forall 1 \leq i \leq m$ if and only if $\xi \equiv j$ on $A_j, j = 0, 1, 2$ for some $A \in \Pi(\Gamma, a, \widehat{\xi}_1, \dots, \widehat{\xi}_m)$. Reasoning as before gives

$$\begin{aligned} & P(\xi_t(x) = a(x) \forall x \in \Gamma, 0 < L^{\Gamma,t} < t_0) \\ &= \sum_{\widehat{\xi}_1, \dots, \widehat{\xi}_m}^{(K)} \sum_{\mathcal{A} \in \Pi(\Gamma, a, \widehat{\xi}_1, \dots, \widehat{\xi}_m)} \int_0^{t_0} P(\widehat{\xi}_s^{x_i} = \widehat{\xi}_i, 1 \leq i \leq m) \times \\ & \quad P\left(\tau_K^{(z,t-s)} = \infty \text{ and } \xi_{t-s}(z) = j \forall z \in A_j, j = 0, 1, 2\right) ds. \end{aligned}$$

By the argument leading to (3.11), for some $\phi(\mathcal{A})$,

$$\begin{aligned} & \lim_{t \rightarrow \infty} P(\xi_t(x) = a(x) \forall x \in \Gamma, 0 < L^{\Gamma,t} \leq t_0) \\ &= \sum_{\widehat{\xi}_1, \dots, \widehat{\xi}_m}^{(K)} \sum_{\mathcal{A} \in \Pi(\Gamma, a, \widehat{\xi}_1, \dots, \widehat{\xi}_m)} \phi(\mathcal{A}) \int_0^{t_0} P(\widehat{\xi}_s^{x_i} = \widehat{\xi}_i, 1 \leq i \leq m) ds. \end{aligned}$$

In view of (3.12), this completes the proof of (3.13). \square

Proof of Theorem 1(iii). We may suppose $\lambda_1 \leq \lambda_2$, $i = 1$ and $j = 2$. Our first task is to prove $\nu_\eta(\xi : |\xi| = \infty) = 1$. To do this it is enough by (ii) to show that for $\varepsilon > 0$ and $M \geq 1$ there is a finite $A \subset S(y)$ such that

$$(3.14) \quad \lim_{t \rightarrow \infty} P\left(\sum_{x \in A} 1\{\xi_t(x) = 1\} \geq M\right) \geq 1 - \varepsilon.$$

To do this, let $\ell > 0$ be large enough so that $(1/\sqrt{d})^\ell \leq \alpha(\lambda_1)/2$. Letting $\varepsilon_0 = \alpha(\lambda_1)/2 > 0$, it follows from (3.3) that for any $y_0 \in S(y)$ and $x_0 \in S(y_0)$ such that $|x_0 - y_0| \geq \ell$,

$$P(\xi_t(x_0) = 1 \text{ and } y_0 \notin D_s^{(x_0, t)} \forall s \leq t) \geq \varepsilon_0 \quad \forall t \geq 0.$$

We construct the set A in (3.14) as follows. Let N be a positive integer large enough so that a binomial random variable X with parameters N and $p \geq \varepsilon_0$ will satisfy $P(X \geq M) \geq 1 - \varepsilon$. Let y_1, \dots, y_N be vertices in $S(y)$ such that $S(y_j) \cap S(y_k) = \emptyset$ for $j \neq k$, and let $x_j \in S(y_j)$ satisfy $|x_j - y_j| \geq \ell$ for $j = 1, \dots, N$. Fix $t > 0$ and define $\varepsilon_j = 1\{\xi_t(x_j) = 1, y_j \notin D_s^{(x_j, t)} \forall s \leq t\}$, $j = 1, \dots, N$. By independence of disjoint space-time regions, the ε_j are independent with $P(\varepsilon_j = 1) \geq \varepsilon_0$, so $X = \sum_{j=1}^N \varepsilon_j$ is binomial with parameters N and $p \geq \varepsilon_0$, and thus (3.14) holds for $A = \{x_i, i = 1, \dots, N\}$.

The fact that $\nu(\xi : \xi(x) = 2) \rightarrow 0$ as $x \rightarrow \infty$, $x \in S(y)$ is a simple consequence of duality and the bound (3.3). Since $x \in S(y)$ and $\xi_0 \equiv 1$ on $S(y)$,

$$P(\xi_t(x) = 2) \leq P(\widehat{\xi}_t^{(x, t)} \cap S^c(y) \neq \emptyset) \leq (1/\sqrt{d})^{|x-y|}.$$

Now let $t \rightarrow \infty$ and $x \rightarrow \infty$, $x \in S(y)$. □

Proof of Theorem 1(iv). Assume $\eta, \eta' \in \Xi_0$. Consider both ξ_t with $\xi_0 = \eta$ and ξ'_t with $\xi'_0 = \eta'$ defined via the Poisson processes in Section 2. Let $A = \{y \in \mathbb{T}_d : \eta(y) \neq \eta'(y)\}$, and suppose A is finite. For any x , since each $\lambda_i \leq \lambda^*$, $P(\widehat{\xi}_t^{(x, t)} \cap A \neq \emptyset) \rightarrow 0$ as $t \rightarrow \infty$. By the duality equation (2.2), since $\xi_0 = \xi'_0$ on A^c , $P(\xi_t(x) \neq \xi'_t(x)) \rightarrow 0$ as $t \rightarrow \infty$, which implies $\nu_\eta = \nu_{\eta'}$. The remaining conclusion of (iv) is a simple consequence of (iii). □

4. PROOF OF THEOREM 2

Before beginning the proof of Theorem 2 we state and prove a fact about the upper invariant measure $\bar{\nu}$ for the single-type contact process. The result we need is an immediate consequence of the inequality I.2.30(b) of [7] for $S = \mathbb{Z}^d$. This inequality also holds for $S = \mathbb{T}_d$, but we could not find a reference. Since the following result is all we need, and it's proof may apply to other choices of S , we give the proof here.

Proposition 1. *Assume S is \mathbb{Z}^d or \mathbb{T}_d and $\bar{\nu}$ is the upper invariant measure for the contact process with birth rate $\lambda > \lambda_*$. Let $\delta_L = \sup_{|A| > L} \bar{\nu}(\zeta : \zeta \cap A = \emptyset)$. Then*

$$(4.1) \quad \delta_L \rightarrow 0 \text{ as } L \rightarrow \infty.$$

Proof. Recall the survival probabilities α_A from (1.8). A consequence of duality for the contact process (see I.1.8 of [6]) is that

$$(4.2) \quad \bar{\nu}(\zeta : \zeta \cap A \neq \emptyset) = \alpha_A.$$

Let ζ_t denote the single-type contact process with birth rate λ . Define the *lifetime* $\sigma(x)$ and the *radius* $r(x)$ of the process started at x by $\sigma(x) = \inf\{t \geq 0 : \zeta_t^x = \emptyset\}$ and $r(x) = \sup\{|y - x| : y \in \zeta_t^x \text{ for some } t \geq 0\}$. Let $\varepsilon_M(x) = P(\sigma(x) < \infty, r(x) \geq M)$.

M). By translation invariance, $\varepsilon_M = \varepsilon_M(x)$ does not depend on x , and $\varepsilon_M \rightarrow 0$ as $M \rightarrow \infty$.

Now fix $M, N > 0$ and let A be any set consisting of N points such that $|x - y| \geq M$ for all $x, y \in A, x \neq y$. Then, by the construction in Section 2,

$$\begin{aligned} 1 - \alpha_A &= P(\sigma(x) < \infty \forall x \in A) \\ &= P(\sigma(x) < \infty, r(x) < M \forall x \in A) + P(\exists x \in A : \sigma(x) < \infty \text{ and } r(x) \geq M). \end{aligned}$$

By independence of disjoint space-time regions and translation invariance, the first probability on the right side equals $\prod_{x \in A} (P(\sigma(x) < \infty, r(x) < M)) \leq (1 - \alpha)^N$ and the second probability is bounded above by $\sum_{x \in A} \varepsilon_M(x) \leq N\varepsilon_M$. Thus

$$1 - \alpha_A \leq (1 - \alpha)^N + N\varepsilon_M.$$

Since $\alpha > 0$, given $\varepsilon > 0$ we may choose first N and then M so that $(1 - \alpha)^N < \varepsilon/2$ and $N\varepsilon_M < \varepsilon/2$. Now let L_0 be large enough so that any $A \subset S$ with at least L_0 points must contain at least N points separated from one another by distance at least M . It follows from monotonicity of α_A that if $|A| > L_0$ then $\bar{\nu}(\zeta : \zeta \cap A = \emptyset) = 1 - \alpha_A < \varepsilon$.

□

Proof of Theorem 2. Here is the idea of the proof. Suppose $t, u > 0$ are large, $\xi_{t+u}(x) \neq 0$ and ${}^2\xi_u \neq \emptyset$. Looking forward from time 0, ${}^2\xi_u$ cannot be empty, and with high probability will have many points. Looking backward from time $t + u$, the dual ancestor process starting at $(x, t + u)$ must survive t time units. For $T > 0$ large but small compared to t , we search for a space-time point (y, s) such that $y \in D_s^{(x, t+u)} = \text{supp}(\tilde{\xi}_s^{(x, t+u)})$ is the primary ancestor at time s , y is 1-blocked, and $D_T^{(y, t+u-s)}$ is nonempty. Trying at most a geometric number of times, with high probability we will find such a point (y, s) with s not too large. Furthermore, $D_T^{(y, t+u-s)} \neq \emptyset$ will imply that $D_{t-s}^{(y, t+u-s)} \neq \emptyset$ with high probability, and also that $D_{t-s}^{(y, t+u-s)}$ will intersect ${}^2\xi_u$. (It is this last point which fails unless $\lambda_2 > \lambda^*$.) This will prevent $\xi_{t+u}(x) = 1$ since the sites in $D_{t-s}^{(y, t+u-s)}$ are descendants of a 1-blocked primary ancestor.

We prepare for the proof of (1.7) by assembling a few preliminary facts. Recall $\bar{\nu} = \bar{\nu}_{\lambda_2}$ from (1.9). Since $\lambda_2 > \lambda^*$, the complete convergence theorem (1.10) implies that for any site x and finite $A \subset S$, with $\alpha = \alpha(\lambda_2)$,

$$(4.3) \quad \lim_{t \rightarrow \infty} P(\zeta_t^x \neq \emptyset, \zeta_t^x \cap A = \emptyset) = \alpha \bar{\nu}(\zeta : \zeta \cap A = \emptyset).$$

Let $\rho(T) = P(\zeta_T^{2,x} \neq \emptyset, \zeta_t^{2,x} = \emptyset \text{ for some } t > T)$, and observe that

$$(4.4) \quad \rho(T) \rightarrow 0 \text{ as } T \rightarrow \infty.$$

Since individuals die at constant rate one, standard arguments show that the 2's must either die out or that their number must tend to infinity, i.e.,

$$(4.5) \quad P(1 \leq |{}^2\xi_u| \leq L) \rightarrow 0 \text{ as } u \rightarrow \infty$$

for fixed $L > 0$.

Our argument uses a certain subset A_t^x of the highest priority ancestors of the forward time ancestor process $\tilde{\xi}_t^x$ which we now define. If $\tilde{\xi}_t^x = \emptyset$ put $A_t^x = \emptyset$. Now

suppose that $\bar{\xi}_t^x = ((a_1(t), b_1(t)), \dots, (a_n(t), b_n(t)))$ for some $n \geq 1$. Put $A_t^x = \emptyset$ if $a_1(t)$ is not 1-blocked (i.e., $b_1(t) = 1$), and otherwise let

$$(4.6) \quad A_t^x = \{a_1(t), \dots, a_m(t)\},$$

where m is the largest index such that $a_1(t), \dots, a_m(t)$ are all 1-blocked. We will see below that for large t , A_t^x is large whenever $\bar{\xi}_t^x \neq \emptyset$.

Our goal is to prove

$$(4.7) \quad \lim_{u \rightarrow \infty} \limsup_{t \rightarrow \infty} P(\xi_{t+u}(x) = 1, |{}^2\xi_u| \geq 1) = 0.$$

This implies (1.7) since $\{\xi_{t+u}(x) = 1, |{}^2\xi_{t+u}| \geq 1\} \subset \{\xi_{t+u}(x) = 1, |{}^2\xi_u| \geq 1\}$. In fact, on account of (4.5), we may focus our attention on $\{\xi_{t+u}(x) = 1, |{}^2\xi_u| > L\}$ for large L .

We begin with an application of the duality equation (2.2),

$$(4.8) \quad P(\xi_{t+u}(x) = 1, |{}^2\xi_u| > L) = P(\Psi(x, \tilde{\xi}_t^{(x, t+u)}, \xi_u) = 1, |{}^2\xi_u| > L).$$

By independence of disjoint space-time regions, and switching to the forward time ancestor process $\bar{\xi}_t^x$ in the second equality,

$$(4.9) \quad \begin{aligned} P(\Psi(x, \tilde{\xi}_t^{(x, t+u)}, \xi_u) = 1, |{}^2\xi_u| > L) &= \int_{|{}^2\eta| > L} P(\xi_u \in d\eta) P(\Psi(x, \tilde{\xi}_t^{(x, t+u)}, \eta) = 1) \\ &= \int_{|{}^2\eta| > L} P(\xi_u \in d\eta) P(\Psi(x, \bar{\xi}_t^x, \eta) = 1). \end{aligned}$$

It follows from the the definitions of Ψ and A_t^x that

$$(4.10) \quad \Psi(x, \bar{\xi}_t^x, \eta) = 2 \text{ on } \{\bar{\xi}_t^x \neq \emptyset, A_t^x \cap {}^2\eta \neq \emptyset\},$$

which implies

$$(4.11) \quad P(\Psi(x, \bar{\xi}_t^x, \eta) = 1) \leq P(\bar{\xi}_t^x \neq \emptyset, A_t^x \cap {}^2\eta = \emptyset).$$

Now combine this with (4.8) and (4.9) to obtain

$$(4.12) \quad P(\xi_{t+u}(x) = 1, |{}^2\xi_u| > L) \leq \int_{|{}^2\eta| > L} P(\xi_u \in d\eta) P(\bar{\xi}_t^x \neq \emptyset, A_t^x \cap {}^2\eta = \emptyset).$$

To show the right side above is small we will argue that for large t and u , $A_t^x \cap {}^2\eta = \emptyset$ is unlikely when $\bar{\xi}_t^x \neq \emptyset$ by means of the following construction. For each $y \in S$ pick some nearest neighbor \tilde{y} , and fix this assignment. Call a space-time point (y, s) *good* if the following events happen:

- (1) a 2-only arrow pointing from \tilde{y} to y occurs during $(s, s+1)$
- (2) a δ occurs at y at some time during $(s+1, s+2)$
- (3) no other events affecting y or \tilde{y} occur during $[s, s+2]$.

Then $\varepsilon_0 = P((y, s) \text{ is good}) > 0$ and does not depend on (y, s) . If $\bar{\xi}_s^x \neq \emptyset$, then $\bar{\xi}_s^x = ((a_1(s), b_1(s)), \dots, (a_n(s), b_n(s)))$ for some $n \geq 1$ and **primary ancestor site** $a_1(s)$. **If $(a_1(s), s)$ is good, then $a_1(s+2) = \tilde{a}_1(s)$, and for $v \geq s+2$ the sites of $\bar{\xi}_v^{(a_1(s+2), s+2)}$ are the highest priority sites of $\bar{\xi}_v^x$, and they are all 1-blocked.**

Fix $T > 0$ and let $s_k = k(T+2)$ and $t_k = s_k + 2$, $k \geq 0$. Let R be the smallest k such that the primary ancestor a in $\bar{\xi}_{s_k}^x$ is good at time s_k and the ancestor process

starting at (\tilde{a}, t_k) lasts at least T time units, i.e.,

$$R = \inf\{k : \bar{\xi}_{s_k}^x \neq \emptyset, (a_1(s_k), s_k) \text{ is good and } \bar{\xi}_{s_{k+1}}^{(a_1(t_k), t_k)} \neq \emptyset\}.$$

Let $\varepsilon_1 = \varepsilon_0 \alpha(\lambda_2) > 0$. By independence of disjoint space-time regions, for any (y, s) and $T > 0$,

$$(4.13) \quad \begin{aligned} P((y, s) \text{ is good and } \bar{\xi}_{s+2+T}^{(\tilde{y}, s+2)} \neq \emptyset) &= P((y, s) \text{ is good})P(\bar{\xi}_{s+2+T}^{(\tilde{y}, s+2)} \neq \emptyset) \\ &= P((y, s) \text{ is good})P(\zeta_{s+2+T}^{2, (\tilde{y}, s+2)} \neq \emptyset) \geq \varepsilon_1. \end{aligned}$$

Iterating this inequality and using the Markov property gives us

$$(4.14) \quad P(\bar{\xi}_{s_k}^x \neq \emptyset \text{ and } R > k) \leq (1 - \varepsilon_1)^k, \quad k \geq 0.$$

Consequently, if $k_0 > 0$ and $t > s_{k_0+1}$,

$$(4.15) \quad P(\bar{\xi}_t^x \neq \emptyset, A_t^x \cap {}^2\eta = \emptyset) \leq (1 - \varepsilon_1)^{k_0} + \sum_{k=0}^{k_0} P(R = k, \bar{\xi}_t^x \neq \emptyset, A_t^x \cap {}^2\eta = \emptyset).$$

Now define the events

$$(4.16) \quad G_k(a) = \{R > k - 1, \bar{\xi}_{s_k}^x \neq \emptyset, a_1(s_k) = a, \text{ and } (a, s_k) \text{ is good}\}, \quad a \in S.$$

For $k \leq k_0$, $\text{supp}(\bar{\xi}_t^{(\tilde{a}, t_k)}) \subset A_t^x$ on $G_k(a)$, and this implies

$$\begin{aligned} P(R = k, \bar{\xi}_t^x \neq \emptyset, A_t^x \cap {}^2\eta = \emptyset) \\ \leq \sum_{a \in S} P(G_k(a) \cap \{\bar{\xi}_{s_{k+1}}^{(\tilde{a}, t_k)} \neq \emptyset, \text{supp}(\bar{\xi}_t^{(\tilde{a}, t_k)}) \cap {}^2\eta = \emptyset\}). \end{aligned}$$

For each a , by independence of disjoint space-time regions,

$$(4.17) \quad \begin{aligned} P(G_k(a) \cap \{\bar{\xi}_{s_{k+1}}^{(\tilde{a}, t_k)} \neq \emptyset, \text{supp}(\bar{\xi}_t^{(\tilde{a}, t_k)}) \cap {}^2\eta = \emptyset\}) \\ \leq P(G_k(a))P(\bar{\xi}_{s_{k+1}}^{(\tilde{a}, t_k)} \neq \emptyset, \text{supp}(\bar{\xi}_t^{(\tilde{a}, t_k)}) \cap {}^2\eta = \emptyset) \\ = P(G_k(a))P(\zeta_T^{2, \tilde{a}} \neq \emptyset, \zeta_{t-t_k}^{2, \tilde{a}} \cap {}^2\eta = \emptyset) \\ \leq P(G_k(a))\left(\rho(T) + P(\zeta_{t-t_k}^{2, \tilde{a}} \neq \emptyset, \zeta_{t-t_k}^{2, \tilde{a}} \cap {}^2\eta = \emptyset)\right), \end{aligned}$$

where we have shifted back to time 0 and used the fact that $t > s_{k_0+1}$. Therefore, by the bounds (4.15) and (4.17), and the fact that $\sum_{a \in S} P(G_k(a)) \leq 1$,

$$(4.18) \quad \begin{aligned} P(\bar{\xi}_t^x \neq \emptyset, A_t^x \cap {}^2\eta = \emptyset) &\leq (1 - \varepsilon_1)^{k_0} + \rho(T)(k_0 + 1) \\ &\quad + \sum_{k=0}^{k_0} \sum_{a \in S} P(G_k(a))P(\zeta_{t-t_k}^{2, \tilde{a}} \neq \emptyset, \zeta_{t-t_k}^{2, \tilde{a}} \cap {}^2\eta = \emptyset). \end{aligned}$$

Recall the definition of δ_L in Proposition 1. For fixed k , a and η such that $|{}^2\eta| > L$, (4.3) implies that

$$\lim_{t \rightarrow \infty} P(\zeta_{t-t_k}^{2, \tilde{a}} \neq \emptyset, \zeta_{t-t_k}^{2, \tilde{a}} \cap {}^2\eta = \emptyset) = \alpha \bar{\nu}(\zeta : \zeta \cap {}^2\eta = \emptyset) \leq \delta_L.$$

With this we can apply Fatou to (4.18) to obtain

$$(4.19) \quad \limsup_{t \rightarrow \infty} P(\bar{\xi}_t^x \neq \emptyset, A_t^x \cap {}^2\eta = \emptyset) \leq (1 - \varepsilon_1)^{k_0} + (k_0 + 1)(\rho(T) + \delta_L),$$

and letting $T \rightarrow \infty$ and using (4.4) then gives us

$$(4.20) \quad \limsup_{t \rightarrow \infty} P(\bar{\xi}_t^x \neq \emptyset, A_t^x \cap {}^2\eta = \emptyset) \leq (1 - \varepsilon_1)^{k_0} + (k_0 + 1)\delta_L.$$

With this inequality we apply Fatou again, this time to (4.12), to obtain

$$(4.21) \quad \limsup_{t \rightarrow \infty} P(\xi_{t+u}(x) = 1, |{}^2\xi_u| > L) \leq (1 - \varepsilon_1)^{k_0} + (k_0 + 1)\delta_L.$$

We have finally established the bound

$$(4.22) \quad \limsup_{t \rightarrow \infty} P(\xi_{t+u}(x) = 1, |{}^2\xi_u| \geq 1) \\ \leq P(1 \leq |{}^2\xi_u| \leq L) + (1 - \varepsilon_1)^{k_0} + (k_0 + 1)\delta_L.$$

Now let $u, L, k_0 \rightarrow \infty$ in order and use (4.5) and (4.1) to finish the proof of (4.7). \square

5. PROOF OF THEOREM 3

Recall the definitions (1.8) and (1.11) and the assumptions of Theorem 3. In view of Theorem 2, it is enough to prove that for all finite $A \subset S$,

$$(5.1) \quad P({}^i\xi_t \cap A \neq \emptyset) \rightarrow \alpha_\eta^i \bar{\nu}_{\lambda_i}(\zeta : \zeta \cap A \neq \emptyset) \text{ as } t \rightarrow \infty, \quad i = 1, 2.$$

By (4.2), $\bar{\nu}_{\lambda_i}(\zeta : \zeta \cap A \neq \emptyset) = \alpha_A(\lambda_i)$, and hence (5.1) is equivalent to

$$(5.2) \quad \lim_{t \rightarrow \infty} P({}^i\xi_t \cap A \neq \emptyset) = \alpha_\eta^i \alpha_A(\lambda_i), \quad i = 1, 2.$$

First consider the case $i = 1$. By (1.7), (5.2) will follow once we establish

$$(5.3) \quad \lim_{t \rightarrow \infty} P({}^1\xi_t \cap A \neq \emptyset, {}^2\xi_t = \emptyset) = \alpha_\eta^1 \alpha_A(\lambda_1).$$

By independence of disjoint space-time regions, for $t, u > 0$,

$$\begin{aligned} P({}^1\xi_{t+u} \cap A \neq \emptyset, {}^2\xi_u = \emptyset) &\leq P({}^1\xi_u \neq \emptyset, {}^2\xi_u = \emptyset, D_t^{1,(x,t+u)} \neq \emptyset \text{ for some } x \in A) \\ &= P({}^1\xi_u \neq \emptyset, {}^2\xi_u = \emptyset) P(D_t^{1,(x,t+u)} \neq \emptyset \text{ for some } x \in A) \\ &= P({}^1\xi_u \neq \emptyset, {}^2\xi_u = \emptyset) P(\zeta_t^{1,A} \neq \emptyset) \\ &\rightarrow \alpha_\eta^1 \alpha_A(\lambda_1) \text{ as } t, u \rightarrow \infty. \end{aligned}$$

Since

$$(5.4) \quad P({}^2\xi_u \neq \emptyset, {}^2\xi_{t+u} = \emptyset) \leq P({}^2\xi_u \neq \emptyset, {}^2\xi_s = \emptyset \text{ for some } s \geq u) \rightarrow 0 \text{ as } u \rightarrow \infty,$$

this proves

$$(5.5) \quad \limsup_{t \rightarrow \infty} P({}^1\xi_t \cap A \neq \emptyset, {}^2\xi_t = \emptyset) \leq \alpha_\eta^1 \alpha_A(\lambda_1).$$

For the required lower bound, duality implies

$$\begin{aligned} P({}^1\xi_{t+u} \cap A \neq \emptyset, {}^2\xi_{t+u} = \emptyset) \\ \geq P({}^2\xi_u = \emptyset, \Psi(x, \widehat{\xi}_t^{(x,t+u)}, \xi_u) = 1 \text{ for some } x \in A). \end{aligned}$$

On the event $\{^2\xi_u = \emptyset\}$, $\Psi(x, \widehat{\xi}_t^{(x,t+u)}, \xi_u) = 1$ if and only if $D_t^{1,(x,t+u)} \cap ^1\xi_u \neq \emptyset$. Consequently, for any $L > 0$,

$$\begin{aligned} P(^1\xi_{t+u} \cap A \neq \emptyset, ^2\xi_{t+u} = \emptyset) &\geq \int_{|^1\eta| > L, ^2\eta = \emptyset} P(\xi_u \in d\eta) P(D_t^{1,(x,t+u)} \cap ^1\eta \neq \emptyset \text{ for some } x \in A) \\ &= \int_{|^1\eta| > L, ^2\eta = \emptyset} P(\xi_u \in d\eta) P(\zeta_t^{1,A} \cap ^1\eta \neq \emptyset). \end{aligned}$$

Now we replace $P(\zeta_t^{1,A} \cap ^1\eta \neq \emptyset)$ with $P(\zeta_t^{1,A} \neq \emptyset) - P(\zeta_t^{1,A} \neq \emptyset, \zeta_t^{1,A} \cap ^1\eta = \emptyset)$ so that the above implies

$$(5.6) \quad P(^1\xi_{t+u} \cap A \neq \emptyset, ^2\xi_{t+u} = \emptyset) \geq P(|^1\xi_u| > L, ^2\xi_u = \emptyset) P(\zeta_t^{1,A} \neq \emptyset) - \int_{|^1\eta| > L, ^2\eta = \emptyset} P(\xi_u \in d\eta) P(\zeta_t^{1,A} \neq \emptyset, \zeta_t^{1,A} \cap ^1\eta = \emptyset).$$

For fixed u and L ,

$$(5.7) \quad \lim_{t \rightarrow \infty} P(|^1\xi_u| > L, ^2\xi_u = \emptyset) P(\zeta_t^{1,A} \neq \emptyset) \geq P(|^1\xi_u| \geq 1, ^2\xi_u = \emptyset) \alpha_A(\lambda_1) - P(1 \leq |^1\xi_u| \leq L).$$

For fixed η with $|^1\eta| > L$, the complete convergence theorem (1.10) implies that

$$(5.8) \quad \lim_{t \rightarrow \infty} P(\zeta_t^{1,A} \neq \emptyset, \zeta_t^{1,A} \cap ^1\eta = \emptyset) = \alpha_A \bar{\nu}_{\lambda_1}(\zeta : \zeta \cap \eta = \emptyset) \leq \delta_L$$

(recall δ_L from Proposition 1. We can now plug (5.7) and (5.8) into (5.6) and use Fatou, keeping u and L fixed, to get

$$(5.9) \quad \liminf_{t \rightarrow \infty} P(^1\xi_{t+u} \cap A \neq \emptyset, ^2\xi_{t+u} = \emptyset) \geq P(|^1\xi_u| \geq 1, ^2\xi_u = \emptyset) \alpha_A(\lambda_1) - P(1 \leq |^1\xi_u| \leq L) - \delta_L.$$

The last two terms on the right side above vanish as first $u \rightarrow \infty$ and then $L \rightarrow \infty$, and therefore $\liminf_{t \rightarrow \infty} P(^1\xi_t \cap A \neq \emptyset, ^2\xi_t = \emptyset) \geq \alpha_\eta^1 \alpha_A(\lambda_1)$. Together with (5.5) this completes the proof of (5.3).

Turning to the $i = 2$ case of (5.2) and using independence of disjoint space-time regions,

$$\begin{aligned} P(^2\xi_{t+u} \cap A \neq \emptyset) &\leq P(^2\xi_u \neq \emptyset, D_t^{(x,t+u)} \neq \emptyset \text{ for some } x \in A) \\ &\leq P(^2\xi_u \neq \emptyset) P(\zeta_t^{2,A} \neq \emptyset). \end{aligned}$$

Now let $t, u \rightarrow \infty$ to obtain

$$(5.10) \quad \limsup_{t \rightarrow \infty} P(^2\xi_t \cap A \neq \emptyset) \leq \alpha_\eta^2 \alpha_A(\lambda_2).$$

For the required lower bound we make use of the 1-blocked ancestor process A_t^x defined in (4.6). By duality and independence of disjoint space-time regions, for any $L > 0$,

$$(5.11) \quad P(^2\xi_{t+u} \cap A \neq \emptyset) \geq \int_{|^2\eta| > L} P(\xi_u \in d\eta) P(\Psi(x, \bar{\xi}_t^x, \eta) = 2 \text{ for some } x \in A).$$

By (4.10),

$$\begin{aligned} P(\Psi(x, \bar{\xi}_t^x, \eta) = 2 \text{ for some } x \in A) &\geq P(\bar{\xi}_t^x \neq \emptyset, A_t^x \cap {}^2\eta \neq \emptyset \text{ for some } x \in A) \\ &\geq P(\bar{\xi}_t^x \neq \emptyset \text{ for some } x \in A) - \sum_{x \in A} P(\bar{\xi}_t^x \neq \emptyset, A_t^x \cap {}^2\eta = \emptyset) \\ &= P(\zeta_t^{2,A} \neq \emptyset) - \sum_{x \in A} P(\bar{\xi}_t^x \neq \emptyset, A_t^x \cap {}^2\eta = \emptyset). \end{aligned}$$

With this bound we can appeal to (4.20), which implies that for any $k_0 > 0$,

$$\begin{aligned} \liminf_{t \rightarrow \infty} P(\Psi(x, \bar{\xi}_t^x, \eta) = 2 \text{ for some } x \in A) \\ \geq \alpha_A(\lambda_2) - |A| \left((1 - \varepsilon_1)^{k_0} + (k_0 + 1)\delta_L \right). \end{aligned}$$

Now we apply Fatou to (5.11) and obtain

$$\begin{aligned} \liminf_{t \rightarrow \infty} P({}^2\xi_{t+u} \cap A \neq \emptyset) &\geq [P({}^2\xi_u \neq \emptyset) - P(1 \leq |{}^2\xi_u| \leq L)] \times \\ &\quad \left[\alpha_A(\lambda_2) - |A| \left((1 - \varepsilon_1)^{k_0} + (k_0 + 1)\delta_L \right) \right]. \end{aligned}$$

Finally, we let $u, L, k_0 \rightarrow \infty$ in order and make use of (4.1) and (4.5) to obtain

$$(5.12) \quad \liminf_{t \rightarrow \infty} P({}^2\xi_t \cap A \neq \emptyset) \geq \alpha_\eta^2 \alpha_A(\lambda_2),$$

which together with (5.10) completes the proof of (5.2) for $i = 2$.

6. PROOF OF THEOREM 4

Recall the notation and definitions of Theorem 4. By (5.10), (5.12) and Theorem 2, it suffices to assume that $|{}^2\eta| < \infty$ and prove

$$(6.1) \quad P({}^1\xi_{t+u} \cap A \neq \emptyset, {}^2\xi_{t+u} = \emptyset) \rightarrow (1 - \alpha_\eta^2)\mu(\zeta : \zeta \cap A \neq \emptyset) \text{ as } t, u \rightarrow \infty.$$

The basic idea of the proof is that when the 2's die out, they “disturb” only a bounded region of space-time. In this case, $\xi_{t+u}(x) = 1$ is essentially the same as $D_{t+u}^{1,(x,t+u)} \cap {}^2\eta \neq \emptyset$, since $\lambda_1 \leq \lambda^*$ implies the 1-dual is unlikely to enter the disturbed space-time region.

Let $\Gamma \subset S$ be a finite set containing ${}^2\eta$ and let ξ_s^Γ be the multitype contact process restricted to Γ . That is, put $\xi_0^\Gamma(x) = \eta(x)$ for $x \in \Gamma$, $\xi_s^\Gamma(x) = 0$ for all $s \geq 0$ and $x \in \Gamma^c$, and let the dynamics of $\xi_s^\Gamma(x)$ for $x \in \Gamma$ be the same as for ξ_s except that only the Poisson processes $T^x, B^{x,y}$, $x, y \in \Gamma$ are used. Consider the events

$$\begin{aligned} E_1 &= \{D_{t+u}^{1,(x,t+u)} \cap {}^1\eta \neq \emptyset \text{ for some } x \in A\}, \\ E_2 &= \{D_s^{1,(x,t+u)} \cap \Gamma = \emptyset \forall x \in A, s \in [t, t+u]\}, \\ E_3 &= \{{}^2\xi_u^\Gamma = \emptyset\}, \\ E_4 &= \{{}^2\xi_s^\Gamma = {}^2\xi_s \forall s \in [0, u]\}. \end{aligned}$$

By independence of disjoint space-time regions, the events $E_1 \cap E_2$ and E_3 are independent, and it is simple to check that

$$(6.2) \quad \bigcap_{i=1}^4 E_i \subset \{{}^1\xi_{t+u} \cap A \neq \emptyset, {}^2\xi_u = \emptyset\} \subset (\bigcap_{i=1}^4 E_i) \cup E_2^c \cup E_4^c.$$

We will prove (6.1) by finding appropriate estimates on the $P(E_i)$ and plugging them into (6.2).

First, by duality,

$$(6.3) \quad P(E_1) = P(\zeta_{t+u}^{1,A} \cap {}^1\eta \neq \emptyset) = P(\zeta_{t+u}^{1,1,\eta} \cap A \neq \emptyset) = P(\zeta_{t+u} \cap A \neq \emptyset).$$

Next, since $\{^2\xi_u = \emptyset\} \subset E_3 \cup E_4^c$ and $E_3 \subset \{^2\xi_u = \emptyset\} \cup E_4^c$,

$$(6.4) \quad P(^2\xi_u = \emptyset) - P(E_4^c) \leq P(E_3) \leq P(^2\xi_u = \emptyset) + P(E_4^c).$$

For fixed finite Γ , switching to forward time, $\lambda_1 \leq \lambda^*$ implies that

$$(6.5) \quad P(E_2^c) \leq \sum_{x \in A} P(\zeta_s^{1,x} \cap \Gamma \neq \emptyset \text{ for some } s \geq t) \rightarrow 0 \text{ as } t \rightarrow \infty,$$

and the fact that ${}^2\eta$ is finite implies that

$$(6.6) \quad P(E_4^c) \rightarrow 0 \text{ as } \Gamma \uparrow S \text{ for fixed } u > 0.$$

For a lower bound on the left side of (6.1), use (6.2), (6.3) and (6.4) to obtain

$$\begin{aligned} P({}^1\xi_{t+u} \cap A \neq \emptyset, {}^2\xi_{t+u} = \emptyset) &\geq P(\cap_{i=1}^3 E_i) - P(E_4^c) \\ &= P(E_1 \cap E_2)P(E_3) - P(E_4^c) \\ &\geq P(E_1)P(E_3) - P(E_2^c) - P(E_4^c) \\ &\geq P(\zeta_{t+u} \cap A \neq \emptyset)P(^2\xi_u = \emptyset) - P(E_2^c) - 2P(E_4^c). \end{aligned}$$

Since $P(\zeta_{t+u} \cap A \neq \emptyset) \rightarrow \mu(\zeta : \zeta \cap A \neq \emptyset)$ as $t \rightarrow \infty$, we let $t \rightarrow \infty$, $\Gamma \uparrow S$, and $u \rightarrow \infty$ in order above and employ (6.5) and (6.6) to obtain

$$\liminf_{t \rightarrow \infty} P(^2\xi_t = \emptyset, {}^1\xi_t \cap A \neq \emptyset) \geq (1 - \alpha_\eta^2)\mu(\zeta : \zeta \cap A \neq \emptyset).$$

By a similar argument,

$$\begin{aligned} P({}^1\xi_{t+u} \cap A \neq \emptyset, {}^2\xi_u = \emptyset) &\leq P(E_1 \cap E_2)P(E_3) + P(E_2^c) + P(E_4^c) \\ &\leq P(\zeta_{t+u} \cap A \neq \emptyset)P(^2\xi_u = \emptyset) + P(E_2^c) + 2P(E_4^c), \end{aligned}$$

and therefore

$$\lim_{u \rightarrow \infty} \limsup_{t \rightarrow \infty} P(^2\xi_{t+u} \cap A \neq \emptyset, {}^1\xi_u = \emptyset) \leq (1 - \alpha_\eta^2)\mu(\zeta : \zeta \cap A \neq \emptyset).$$

Combining this with (5.4)

$$P(^2\xi_u \neq \emptyset, {}^2\xi_{t+u} = \emptyset) \leq P(^2\xi_u \neq \emptyset, {}^2\xi_s = \emptyset \text{ for some } s \geq u) \rightarrow 0 \text{ as } u \rightarrow \infty,$$

gives us

$$\limsup_{t \rightarrow \infty} P(^2\xi_t = \emptyset, {}^1\xi_t \cap A \neq \emptyset) \leq (1 - \alpha_\eta^2)\mu(\zeta : \zeta \cap A \neq \emptyset),$$

and we are done.

ACKNOWLEDGMENTS

We thank Rick Durrett for showing us the proof of (1.7) and Robin Pemantle for the argument for (4.1).

References.

- [1] B. Chan and R. Durrett (2006). A new coexistence result for competing contact processes. *Ann. Appl. Probab.*, **16**, 1155-1165.
- [2] R. Durrett and C. Neuhauser (1997). Coexistence results for some models. *Ann. Appl. Probab.* **7**, 10-45.
- [3] O. Haggstrom and R. Pemantle (2000). Absence of mutual unbounded growth for almost all parameter values in the two-type Richardson model. *Stochastic Process. Appl.*, **90**, 207-222.
- [4] T.E.Harris (1974). Contact interactions on a lattice. *Ann. Probab.* **2**, 969-988.
- [5] G. Kordzakhia and S. Lalley (2005). A two species competition model on \mathbb{Z}^d . *Stochastic Process. Appl.*, **115**, 781-796.
- [6] T. Liggett (1985). *Interacting particle systems*, Springer-Verlag.
- [7] T.Liggett (1999). *Stochastic interacting systems: contact, voter and exclusion processes*, Springer-Verlag, Berlin.
- [8] C. Neuhauser (1992). Ergodic theorems for the multitype contact process. *Probab. Theory Related Fields*, **91**, 467-506.

MATHEMATICS DEPARTMENT, SYRACUSE UNIVERSITY, SYRACUSE, NY 13244
E-mail address: `jtc Cox@syr.edu`

MATHEMATICS DEPARTMENT, UNIVERSITY OF COLORADO, COLORADO SPRINGS, CO 80933
E-mail address: `rschinaz@uccs.edu`